



CTU

CZECH TECHNICAL
UNIVERSITY
IN PRAGUE

Electron in triangular graphene dots

J. Hrivnák, L. Motlochová

Czech Technical University in Prague, Faculty of Nuclear Sciences and
Physical Engineering, Břehová 7, 115 19 Praha 1

September 6, 2022

AAMP XIX, Prague, Czech Republic

Motivation

General aim

(non-relativistic) electron propagation in triangular graphene quantum dots

Mathematical part

- honeycomb lattice via the root and weight lattices of the crystallographic root system A_2
- armchair and zigzag honeycomb Fourier–Weyl transforms

Physical part

- Hamiltonian description of discrete quantum billiard-type systems on honeycomb lattice
- incorporation of boundary conditions



References

- J. Hrivnák, L. Motlochová, *On electron propagation in triangular graphene quantum dots*, J. Phys. A: Math. Theor. **55** (2022) 125201
- J. Hrivnák, L. Motlochová, *Discrete cosine and sine transforms generalized to honeycomb lattice*, J. Math. Phys. **59** (2018) 063503
- J. Hrivnák, L. Motlochová, *Discrete cosine and sine transforms generalized to honeycomb lattice II. Zigzag boundaries*, J. Math. Phys. **62** (2021) 093504
- T. Czyżycki, J. Hrivnák, L. Motlochová, *Generalized Dual-Root Lattice Transforms of Affine Weyl Groups*, Symmetry **12** (2020) 1018
- A. Brus, J. Hrivnák, L. Motlochová, *Quantum particle on dual root lattice in Weyl alcove*, J. Phys. A: Math. Theor. **54** (2021) 095202
- A. U. Klimyk, J. Patera, *Orbit functions*, SIGMA **2** (2006) 006
- R. V. Moody, J. Patera, *Orthogonality within the families of C -, S -, and E -functions of any compact semisimple Lie group*, SIGMA **2** (2006) 076



Outline

1 Honeycomb Fourier–Weyl Transforms

- Root and weight lattices
- Triangular honeycomb dots
- Honeycomb Weyl orbit functions

2 Electron propagation in TGQDs

- Hamiltonians
- Time-independent Schrödinger equations
- Electronic probability densities

Root system A_2

- the **irreducible crystallographic root system** $\Pi(A_2)$ of rank 2
- the set of **simple roots** $\Delta = \{\alpha_1, \alpha_2\} \subset \Pi$, $\text{span}_{\mathbb{R}} \Delta = \mathbb{R}^2$, scalar product $\langle \cdot, \cdot \rangle$
- roots of the same length with relative angle $2\pi/3$

$$\langle \alpha_1, \alpha_1 \rangle = \langle \alpha_2, \alpha_2 \rangle = 2, \quad \langle \alpha_1, \alpha_2 \rangle = -1$$

- \mathbb{Z} -dual ω -basis $\{\omega_1, \omega_2\}$

$$\langle \omega_i, \alpha_j \rangle = \delta_{ij}, \quad i, j \in \{1, 2\}$$

- reflections** r_i , $i \in \{1, 2\}$ for $x \in \mathbb{R}^2$

$$r_i x = x - \langle x, \alpha_i \rangle \alpha_i$$

- Weyl group** W generated by r_1, r_2
- group of the **sign homomorphisms** $\Sigma = \{1, \sigma^e\} \simeq \mathbb{Z}_2$

$$1(w) = 1, \quad \sigma^e(w) = \det(w)$$



Root and weight lattices

- **even Weyl group** W^e

$$W^e = \{w \in W \mid \sigma^e(w) = 1\}$$

- **root lattice** Q and **weight lattice** P

$$Q = \mathbb{Z}\alpha_1 + \mathbb{Z}\alpha_2$$

$$P = \mathbb{Z}\omega_1 + \mathbb{Z}\omega_2$$

- disjoint **decompositions of the lattices** P and $\frac{1}{3}Q$

$$P = Q \cup \{\omega_1 + Q\} \cup \{-\omega_1 + Q\}$$

$$\frac{1}{3}Q = P \cup \left\{\frac{1}{3}\alpha_1 + P\right\} \cup \left\{-\frac{1}{3}\alpha_1 + P\right\}$$

- **extended affine Weyl group** W_P^{aff} and **affine Weyl group** $W^{\text{aff}} \subset W_P^{\text{aff}}$

$$W_P^{\text{aff}} = P \rtimes W$$

$$W^{\text{aff}} = Q \rtimes W$$

- **fundamental domains** $F_P \subset F \subset \mathbb{R}^2$
- coordinate x_0 of $x = x_1\omega_1 + x_2\omega_2 \in MF$, $x_0 = M - x_1 - x_2$

$$x \equiv [x_0, x_1, x_2]_M$$

Counting functions

- magnifying factor $M \in \mathbb{N}$, **counting functions** $h_M : MF \rightarrow \mathbb{N}$ and $\varepsilon : F \rightarrow \mathbb{N}$

$$h_M(x) = |\text{Stab}_{W^{\text{aff}}} \left(\frac{x}{M} \right)|, \quad \varepsilon(x) = \frac{|W|}{h_1(x)}$$

- **retraction homomorphism** $\psi : W^{\text{aff}} \rightarrow W$ for $z = T(q)w \in W^{\text{aff}}$

$$\psi(z) = w$$

- for any $a \in \mathbb{R}^2$ there exists $a' \in F$ and $z[a] \in W^{\text{aff}}$ such that

$$a = z[a]a'$$

Function $\chi^\sigma : \mathbb{R}^2 \rightarrow \{-1, 0, 1\}$

$$\chi^\sigma(a) = \begin{cases} \sigma \circ \psi(z[a]), & \sigma \circ \psi(\text{Stab}_{W^{\text{aff}}}(a)) = 1 \\ 0, & \sigma \circ \psi(\text{Stab}_{W^{\text{aff}}}(a)) = \{1, -1\} \end{cases}$$

- **signed fundamental domain** $F^\sigma \subset F$

$$F^\sigma = \{x \in F \mid \chi^\sigma(x) = 1\}$$

Outline

1 Honeycomb Fourier–Weyl Transforms

- Root and weight lattices
- **Triangular honeycomb dots**
- Honeycomb Weyl orbit functions

2 Electron propagation in TGQDs

- Hamiltonians
- Time-independent Schrödinger equations
- Electronic probability densities

Armchair honeycomb dots

- W^{aff} -invariant **honeycomb lattice** A_M subtractively

$$A_M = \frac{1}{M}(P \setminus Q) = A_M^+ \cup A_M^-$$

- sublattices** of the honeycomb lattice A_M

$$A_M^+ = \frac{1}{M}(\omega_1 + Q)$$

$$A_M^- = \frac{1}{M}(-\omega_1 + Q)$$

- armchair sign function** $\text{sga} : A_M \rightarrow \{1, -1\}$, $\text{sga}(wa) = \text{sga}(a)$, $w \in W$

$$\text{sga}(a) = \begin{cases} 1, & a \in A_M^+, \\ -1, & a \in A_M^- \end{cases}$$

- armchair neighbourhood sets** $B_{A,M}^{(k)}(a) \subset A_M$, $k \in \{0, 1, 2\}$

$$B_{A,M}^{(0)}(a) = a$$

$$B_{A,M}^{(1)}(a) = a + \text{sga}(a) W^e \left(\frac{\omega_1}{M} \right)$$


$$B_{A,M}^{(2)}(a) = a + W \left(\frac{\alpha_1}{M} \right)$$

- armchair honeycomb dots** $H_{A,M}^\sigma$, $\sigma \in \Sigma$

$$H_{A,M}^\sigma = A_M \cap F^\sigma$$



Armchair honeycomb dots $H_{A,6}^\sigma$



Armchair weight sets

- weight sets $L_M^\sigma \subset P, \sigma \in \Sigma$

$$L_M^\sigma = P \cap MF^\sigma$$

- weight sets in Kac coordinates

$$L_M^1 = \left\{ [\lambda_0, \lambda_1, \lambda_2]_M \mid \lambda_0, \lambda_1, \lambda_2 \in \mathbb{Z}^{\geq 0} \right\}$$

$$L_M^{\sigma^e} = \{ [\lambda_0, \lambda_1, \lambda_2]_M \mid \lambda_0, \lambda_1, \lambda_2 \in \mathbb{N} \}$$

- armchair weight sets** $L_{A,M}^{\sigma,\pm}$

$$L_{A,M}^{\sigma,\pm} = \{ \lambda \in L_M^\sigma \mid (\lambda_0 > \lambda_1, \lambda_0 > \lambda_2) \vee (\lambda_0 = \lambda_1 > \lambda_2) \}$$



- sums of numbers of weights in $L_{A,M}^{\sigma,+}$ and $L_{A,M}^{\sigma,-}$

$$\left| L_{A,M}^{\sigma,+} \right| + \left| L_{A,M}^{\sigma,-} \right| = \left| H_{A,M}^\sigma \right|$$

- armchair weight transformation** $\gamma_2 : L_M^1 \rightarrow P$ for $\lambda \in L_M^1$

$$\gamma_2 \lambda = (r_1 r_2)^2 \lambda + M \omega_2$$

Armchair honeycomb dots $H_{A,7}^\sigma$ and weight sets $L_{A,7}^{\sigma,\pm}$



Zigzag honeycomb dots

- W^{aff} -invariant **honeycomb lattice** Z_M subtractively

$$Z_M = \frac{1}{M}(\frac{1}{3}Q \setminus P) = Z_M^+ \cup Z_M^-$$

- **sublattices** of the honeycomb lattice Z_M

$$Z_M^+ = \frac{1}{M}(\frac{1}{3}\alpha_1 + P)$$

$$Z_M^- = \frac{1}{M}(-\frac{1}{3}\alpha_1 + P)$$

- **zigzag sign function** $\text{sgz} : Z_M \rightarrow \{1, -1\}$, $\text{sgz}(wa) = \sigma^e(w) \text{sgz}(a)$,
 $w \in W$

$$\text{sgz}(a) = \begin{cases} 1, & a \in Z_M^+, \\ -1, & a \in Z_M^- \end{cases}$$

- **zigzag neighbourhood sets** $B_{Z,M}^{(k)}(a) \subset Z_M$, $k \in \{0, 1, 2\}$

$$B_{Z,M}^{(0)}(a) = a$$

$$B_{Z,M}^{(1)}(a) = a + \text{sgz}(a) W^e \left(\frac{\alpha_1}{3M} \right)$$


$$B_{Z,M}^{(2)}(a) = a + \left[W^e \left(\frac{\omega_1}{M} \right) \cup W^e \left(-\frac{\omega_1}{M} \right) \right]$$

- **zigzag honeycomb dots** $H_{Z,M}^\sigma$, $\sigma \in \Sigma$

$$H_{Z,M}^\sigma = Z_M \cap F^\sigma$$



Zigzag honeycomb dots $H_{Z,4}^\sigma$



Zigzag weight sets

- zigzag boundary weight subsets of L_M^1

$$L_M^{Z,a} = \left\{ [0, \lambda_1, \lambda_2]_M \mid \lambda_1, \lambda_2 \in \mathbb{N} \right\}$$

$$L_M^{Z,c} = \left\{ [\lambda_0, 0, \lambda_2]_M \mid \lambda_0, \lambda_2 \in \mathbb{N} \right\} \cup \left\{ [\lambda_0, \lambda_1, 0]_M \mid \lambda_0, \lambda_1 \in \mathbb{N} \right\}$$

$$O_M^Z = \{[M, 0, 0]_M\}$$

- **zigzag weight sets** $L_{Z,M}^{\sigma,\pm}$

$$L_{Z,M}^{1,+} = L_{Z,M}^{\sigma^e,-} = L_M^{\sigma^e} \cup L_M^{Z,a}$$

$$L_{Z,M}^{1,-} = L_{Z,M}^{\sigma^e,+} = L_M^{\sigma^e} \cup O_M^Z \cup L_M^{Z,c}$$



- sums of numbers of weights in $L_{Z,M}^{\sigma,+}$ and $L_{Z,M}^{\sigma,-}$

$$|L_{Z,M}^{\sigma,+}| + |L_{Z,M}^{\sigma,-}| = |H_{Z,M}^{\sigma}|$$

- **zigzag weight transformation** $\beta_1 : L_M^1 \rightarrow P$ for $\lambda \in L_M^1$

$$\beta_1 \lambda = r_1 r_2 r_1 \lambda + M(\omega_1 + \omega_2)$$

Zigzag honeycomb dots $H_{Z,4}^\sigma$ and weight sets $L_{Z,4}^{\sigma,\pm}$



Outline

1 Honeycomb Fourier–Weyl Transforms

- Root and weight lattices
- Triangular honeycomb dots
- Honeycomb Weyl orbit functions

2 Electron propagation in TGQDs

- Hamiltonians
- Time-independent Schrödinger equations
- Electronic probability densities



Weyl orbit functions

- **Weyl orbit functions** (C - and S -functions) $\varphi_b^\sigma : \mathbb{R}^2 \rightarrow \mathbb{C}, \sigma \in \Sigma, b \in \mathbb{R}^2, x \in \mathbb{R}^2$

$$\varphi_b^\sigma(x) = \sum_{w \in W} \sigma(w) e^{2\pi i \langle wb, x \rangle}$$

- **even Weyl orbit functions** (E -functions) $\Xi_b : \mathbb{R}^2 \rightarrow \mathbb{C}, b \in \mathbb{R}^2, x \in \mathbb{R}^2$

$$\Xi_b(x) = \sum_{w \in W^e} e^{2\pi i \langle wb, x \rangle}$$

- product-to-sum formulas of φ_b^σ and $\varphi_b^{\sigma'}, \sigma, \sigma' \in \Sigma, x, y \in \mathbb{R}^2$

$$\varphi_b^\sigma(x) \varphi_b^{\sigma'}(y) = \sum_{w \in W} \sigma'(w) \varphi_b^{\sigma \cdot \sigma'}(x + wy)$$

- (anti)symmetries for $x' \in F$ and $x \in W^{\text{aff}} x'$

$$\varphi_\lambda^\sigma(x) = \chi^\sigma(x) \varphi_\lambda^\sigma(x')$$

Armchair orbit functions

- extension coefficients $\mu_{0,\lambda}^{A,\sigma,\pm}$ and $\mu_{2,\lambda}^{A,\sigma,\pm}$, $\sigma \in \Sigma$, $\lambda \in L_{A,M}^{\sigma,\pm}$

$$\mu_{0,\lambda}^{A,\sigma,\pm} = \operatorname{Re} \left\{ \left(1 + \frac{i}{\sqrt{3}} \right) \Xi_{\lambda} \left(\frac{\omega_1}{M} \right) \right\}$$

$$\mu_{2,\lambda}^{A,\sigma,\pm} = \operatorname{Re} \left\{ \left(1 - \frac{i}{\sqrt{3}} \right) \Xi_{\lambda} \left(\frac{\omega_1}{M} \right) \right\} \pm \left| \Xi_{\lambda} \left(\frac{\omega_1}{M} \right) \right|$$

- armchair orbit functions** $h_{\lambda}^{A,\sigma,\pm} : \mathbb{R}^2 \rightarrow \mathbb{C}$, $\lambda \in L_{A,M}^{\sigma,\pm}$, $\sigma \in \Sigma$, $x \in \mathbb{R}^2$

$$h_{\lambda}^{A,\sigma,\pm}(x) = \mu_{0,\lambda}^{A,\sigma,\pm} \varphi_{\lambda}^{\sigma}(x) + \mu_{2,\lambda}^{A,\sigma,\pm} \varphi_{\gamma_2 \lambda}^{\sigma}(x)$$

- normalization functions $\mu^{A,\sigma,\pm} : L_{A,M}^{\sigma,\pm} \rightarrow \mathbb{R}^{>0}$ for $\lambda \in L_{A,M}^{\sigma,\pm}$

$$\mu^{A,\sigma,\pm}(\lambda) = \left| \Xi_{\lambda} \left(\frac{\omega_1}{M} \right) \right| \left(2 \left| \Xi_{\lambda} \left(\frac{\omega_1}{M} \right) \right| \pm \operatorname{Re} \left\{ (1 - \sqrt{3}i) \Xi_{\lambda} \left(\frac{\omega_1}{M} \right) \right\} \right)$$

- discrete orthogonality relations**

$$\sum_{a \in H_{A,M}^{\sigma}} \varepsilon(a) \left[h_{\lambda}^{A,\sigma,\pm}(a) \right]^* h_{\lambda'}^{A,\sigma,\pm}(a) = 12M^2 h_M(\lambda) \mu^{A,\sigma,\pm}(\lambda) \delta_{\lambda\lambda'}, \quad \lambda, \lambda' \in L_{A,M}^{\sigma,\pm}$$

$$\sum_{a \in H_{A,M}^{\sigma}} \varepsilon(a) \left[h_{\lambda}^{A,\sigma,+}(a) \right]^* h_{\lambda'}^{A,\sigma,-}(a) = 0, \quad \lambda \in L_{A,M}^{\sigma,+}, \quad \lambda' \in L_{A,M}^{\sigma,-}$$



Zigzag orbit functions

- extension coefficients $\mu_{0,\lambda}^{Z,\sigma,\pm}$ and $\mu_{1,\lambda}^{Z,\sigma,\pm}$, $\sigma \in \Sigma$, $\lambda \in L_{Z,M}^{\sigma,\pm}$

$$\mu_{0,\lambda}^{Z,\sigma,\pm} = \sigma(r_1) \operatorname{Re} \left\{ \left(1 - \frac{i}{\sqrt{3}} \right) \Xi_\lambda \left(\frac{\alpha_1}{3M} \right) \right\}$$

$$\mu_{1,\lambda}^{Z,\sigma,\pm} = \operatorname{Re} \left\{ \left(1 + \frac{i}{\sqrt{3}} \right) \Xi_\lambda \left(\frac{\alpha_1}{3M} \right) \right\} \pm \left| \Xi_\lambda \left(\frac{\alpha_1}{3M} \right) \right|$$

- zigzag orbit functions** $h_\lambda^{Z,\sigma,\pm} : \mathbb{R}^2 \rightarrow \mathbb{C}$, $\lambda \in L_{Z,M}^{\sigma,\pm}$, $\sigma \in \Sigma$, $x \in \mathbb{R}^2$

$$h_\lambda^{Z,\sigma,\pm}(x) = \mu_{0,\lambda}^{Z,\sigma,\pm} \varphi_\lambda^\sigma(x) + \mu_{1,\lambda}^{Z,\sigma,\pm} \varphi_{\beta_1 \lambda}^\sigma(x).$$

- normalization functions $\mu^{Z,\sigma,\pm} : L_{Z,M}^{\sigma,\pm} \rightarrow \mathbb{R}^{>0}$ for $\lambda \in L_{Z,M}^{\sigma,\pm}$

$$\mu^{Z,\sigma,\pm}(\lambda) = 2 \left| \Xi_\lambda \left(\frac{\alpha_1}{3M} \right) \right| \left(2 \left| \Xi_\lambda \left(\frac{\alpha_1}{3M} \right) \right| \pm \operatorname{Re} \left\{ \left(1 + \sqrt{3}i \right) \Xi_\lambda \left(\frac{\alpha_1}{3M} \right) \right\} \right)$$

- discrete orthogonality relations**

$$\sum_{a \in H_{Z,M}^\sigma} \left[h_\lambda^{Z,\sigma,\pm}(a) \right]^* h_{\lambda'}^{Z,\sigma,\pm}(a) = 3M^2 \mu^{Z,\sigma,\pm}(\lambda) \delta_{\lambda\lambda'}, \quad \lambda, \lambda' \in L_{Z,M}^{\sigma,\pm},$$

$$\sum_{a \in H_{Z,M}^\sigma} \left[h_\lambda^{Z,\sigma,+}(a) \right]^* h_{\lambda'}^{Z,\sigma,-}(a) = 0, \quad \lambda \in L_{Z,M}^{\sigma,+}, \lambda' \in L_{Z,M}^{\sigma,-}$$

Outline

- 1 Honeycomb Fourier–Weyl Transforms
 - Root and weight lattices
 - Triangular honeycomb dots
 - Honeycomb Weyl orbit functions

- 2 Electron propagation in TGQDs
 - **Hamiltonians**
 - Time-independent Schrödinger equations
 - Electronic probability densities

Armchair Hamiltonians

- **energy parameters** $I_k \in \mathbb{R}$, with $k \in \{0, 1, 2\}$ and I_1 positive, an electron with the on-site energy $E_0 = -I_0$ propagating with amplitudes iI_1/\hbar and iI_2/\hbar per unit time to the nearest and next-to-nearest lattice positions
- orthonormal **armchair position bases** $|a; A, \sigma\rangle$, $\sigma \in \Sigma$, $a \in H_{A,M}^\sigma$ span the complex Hilbert spaces $\mathcal{H}_{A,M}^\sigma$

- coupling sets

$$N_{A,M}^{(k)}(a, a') = W^{\text{aff}} a' \cap B_{A,M}^{(k)}(a)$$

- **hopping operators** for $a, a' \in H_{A,M}^\sigma$

$$\left\langle a; A, \sigma \left| \widehat{I}_{A,M}^{\sigma, (k)} \right| a'; A, \sigma \right\rangle = -I_k \varepsilon^{\frac{1}{2}}(a) \varepsilon^{-\frac{1}{2}}(a') \sum_{d \in N_{A,M}^{(k)}(a, a')} \chi^\sigma(d)$$

- **tight-binding Hamiltonians** $\widehat{H}_{A,M}^\sigma : \mathcal{H}_{A,M}^\sigma \rightarrow \mathcal{H}_{A,M}^\sigma$

$$\widehat{H}_{A,M}^\sigma = \widehat{I}_{A,M}^{\sigma, (0)} + \widehat{I}_{A,M}^{\sigma, (1)} + \widehat{I}_{A,M}^{\sigma, (2)}$$

Zigzag Hamiltonians

- orthonormal **zigzag position bases** $|a; Z, \sigma\rangle, \sigma \in \Sigma, a \in H_{Z,M}^\sigma$ span the complex Hilbert spaces $\mathcal{H}_{Z,M}^\sigma$

- coupling sets

$$N_{Z,M}^{(k)}(a, a') = W^{\text{aff}} a' \cap B_{Z,M}^{(k)}(a)$$

- **hopping operators** for $a, a' \in H_{Z,M}^\sigma$

$$\left\langle a; Z, \sigma \left| \hat{I}_{Z,M}^{\sigma, (k)} \right| a'; Z, \sigma \right\rangle = -I_k \sum_{d \in N_{Z,M}^{(k)}(a, a')} \chi^\sigma(d)$$

- **tight-binding Hamiltonians** $\hat{H}_{Z,M}^\sigma : \mathcal{H}_{Z,M}^\sigma \rightarrow \mathcal{H}_{Z,M}^\sigma$

$$\hat{H}_{Z,M}^\sigma = \hat{I}_{Z,M}^{\sigma, (0)} + \hat{I}_{Z,M}^{\sigma, (1)} + \hat{I}_{Z,M}^{\sigma, (2)}$$

Armchair momentum bases

- armchair orthonormal **momentum bases** $|\lambda; A, \sigma, \pm\rangle \in \mathcal{H}_{A,M}^\sigma$,
 $\lambda \in L_{A,M}^{\sigma,\pm}$

$$|\lambda; A, \sigma, \pm\rangle = \sum_{a \in H_{A,M}^\sigma} |a; A, \sigma\rangle \langle a; A, \sigma | \lambda; A, \sigma, \pm\rangle$$

where

$$\langle a; A, \sigma | \lambda; A, \sigma, \pm\rangle = \sqrt{\frac{\varepsilon(a)}{12M^2 h_M(\lambda) \mu^{A,\sigma,\pm}(\lambda)}} h_\lambda^{A,\sigma,\pm}(a)$$

- energy functions** $E_A^{(1),\pm}, E_A^{(2)} : F_P \rightarrow \mathbb{R}$ for $x \in F_P$

$$E_A^{(1),\pm}(x) = \pm I_1 |\Xi_{\omega_1}(x)|$$

$$E_A^{(2)}(x) = -2I_2 \operatorname{Re} \{\Xi_{\alpha_1}(x)\}$$

Zigzag momentum bases

- zigzag orthonormal **momentum bases** $|\lambda; Z, \sigma, \pm\rangle \in \mathcal{H}_{Z,M}^\sigma$, $\lambda \in L_{Z,M}^{\sigma,\pm}$

$$|\lambda; Z, \sigma, \pm\rangle = \sum_{a \in H_{Z,M}^\sigma} |a; Z, \sigma\rangle \langle a; Z, \sigma | \lambda; Z, \sigma, \pm\rangle,$$

where

$$\langle a; Z, \sigma | \lambda; Z, \sigma, \pm\rangle = \sqrt{\frac{1}{3M^2 \mu^{Z,\sigma,\pm}(\lambda)}} h_\lambda^{Z,\sigma,\pm}(a)$$

- energy functions** $E_Z^{(1),\pm}, E_Z^{(2)} : F \rightarrow \mathbb{R}$ for $x \in F$

$$E_Z^{(1),\pm}(x) = \pm I_1 \left| \Xi_{\frac{\alpha_1}{3}}(x) \right|$$

$$E_Z^{(2)}(x) = -2I_2 \operatorname{Re} \{ \Xi_{\omega_1}(x) \}$$

Outline

- 1 Honeycomb Fourier–Weyl Transforms
 - Root and weight lattices
 - Triangular honeycomb dots
 - Honeycomb Weyl orbit functions

- 2 Electron propagation in TGQDs
 - Hamiltonians
 - Time-independent Schrödinger equations
 - Electronic probability densities

Time-independent armchair Schrödinger equations

Theorem

Let $L_{A,M}^{\sigma,\pm}$ be the armchair weight sets which correspond to the magnifying factor $M \in \mathbb{N}$ and sign homomorphism $\sigma \in \Sigma$ and let $|\lambda; A, \sigma, \pm\rangle \in \mathcal{H}_{A,M}^{\sigma}$, $\lambda \in L_{A,M}^{\sigma,\pm}$ be the vectors of the armchair momentum basis. Then **each vector** $|\lambda; A, \sigma, \pm\rangle \in \mathcal{H}_{A,M}^{\sigma}$ **satisfies** for the armchair Hamiltonian $\hat{H}_{A,M}^{\sigma}$ the **time-independent armchair Schrödinger equation**,

$$\hat{H}_{A,M}^{\sigma} |\lambda; A, \sigma, \pm\rangle = E_{A,\lambda,M}^{\sigma,\pm} |\lambda; A, \sigma, \pm\rangle,$$

with the **eigenenergy** $E_{A,\lambda,M}^{\sigma,\pm} \in \mathbb{R}$ determined by the armchair energy functions as

$$E_{A,\lambda,M}^{\sigma,\pm} = E_0 + E_A^{(1),\pm} \left(\frac{\lambda}{M} \right) + E_A^{(2)} \left(\frac{\lambda}{M} \right).$$



Time-independent zigzag Schrödinger equations

Theorem

Let $L_{Z,M}^{\sigma,\pm}$ be the zigzag weight sets which correspond to the magnifying factor $M \in \mathbb{N}$ and sign homomorphism $\sigma \in \Sigma$ and let $|\lambda; Z, \sigma, \pm\rangle \in \mathcal{H}_{Z,M}^{\sigma}$, $\lambda \in L_{Z,M}^{\sigma,\pm}$ be the vectors of the zigzag momentum basis. Then **each vector** $|\lambda; Z, \sigma, \pm\rangle \in \mathcal{H}_{Z,M}^{\sigma}$ **satisfies** for the zigzag Hamiltonian $\hat{H}_{Z,M}^{\sigma}$ the **time-independent zigzag Schrödinger equation**,



$$\hat{H}_{Z,M}^{\sigma} |\lambda; Z, \sigma, \pm\rangle = E_{Z,\lambda,M}^{\sigma,\pm} |\lambda; Z, \sigma, \pm\rangle,$$

with the **eigenenergy** $E_{Z,\lambda,M}^{\sigma,\pm} \in \mathbb{R}$ determined by the zigzag energy functions as



$$E_{Z,\lambda,M}^{\sigma,\pm} = E_0 + E_Z^{(1),\pm} \left(\frac{\lambda}{M} \right) + E_Z^{(2)} \left(\frac{\lambda}{M} \right).$$



Electronic band structure of the triangular armchair graphene dots



Electronic band structure of the triangular zigzag graphene dots



Outline

- 1 Honeycomb Fourier–Weyl Transforms
 - Root and weight lattices
 - Triangular honeycomb dots
 - Honeycomb Weyl orbit functions

- 2 Electron propagation in TGQDs
 - Hamiltonians
 - Time-independent Schrödinger equations
 - Electronic probability densities



Armchair electronic probability densities

Theorem

Let $L_{A,M}^{\sigma,\pm}$ be the armchair weight sets which correspond to the magnifying factor $M \in \mathbb{N}$ and sign homomorphism $\sigma \in \Sigma$ and let $P_M[\lambda; A, \sigma, \pm]$, $\lambda \in L_{A,M}^{\sigma,\pm}$ be the armchair electronic probability densities. Then the **armchair probability densities** $P_M[\lambda; A, \sigma, +]$ and $P_M[\lambda; A, \sigma, -]$ **coincide** for each $\lambda \in L_{A,M}^{\sigma,+}$ and evaluate for $a \in H_{A,M}^\sigma$ as

$$P_M[\lambda; A, \sigma, \pm](a) = |\langle a; A, \sigma \mid \lambda; A, \sigma, \pm \rangle|^2 = \frac{\varepsilon(a)}{12M^2 h_M(\lambda)} |\varphi_\lambda^\sigma(a)|^2.$$



Zigzag electronic probability densities

Theorem

Let $L_{Z,M}^{\sigma,\pm}$ be the zigzag weight sets which correspond to the magnifying factor $M \in \mathbb{N}$ and sign homomorphism $\sigma \in \Sigma$ and let $P_M[\lambda; Z, \sigma, \pm]$, $\lambda \in L_{Z,M}^{\sigma,\pm}$ be the zigzag electronic probability densities given for $a \in H_{Z,M}^{\sigma}$ as


$$P_M[\lambda; Z, \sigma, \pm](a) = |\langle a; Z, \sigma \mid \lambda; Z, \sigma, \pm \rangle|^2 = \frac{1}{3M^2 \mu^{Z,\sigma,\pm}(\lambda)} \left| h_{\lambda}^{Z,\sigma,\pm}(a) \right|^2.$$

Then the **zigzag probability densities** $P_M[\lambda; Z, \sigma^e, \pm]$ **and** $P_M[\lambda; Z, 1, \mp]$ **coincide** for any $\lambda \in L_{Z,M}^{\sigma^e,\pm}$,


$$P_M[\lambda; Z, \sigma^e, \pm] = P_M[\lambda; Z, 1, \mp].$$

Electronic probability densities of the triangular armchair dots


$P_{50}[(1, 1); A, \sigma^e, \pm]$




$P_{50}[(2, 1); A, \sigma^e, \pm]$




$P_{50}[(3, 1); A, \sigma^e, \pm]$




$P_{50}[(1, 1); A, 1, \pm]$



$P_{50}[(2, 1); A, 1, \pm]$




$P_{50}[(3, 1); A, 1, \pm]$




Electronic probability densities of the triangular zigzag dots


$P_{29}[(1, 1); Z, \sigma^e, -]$




$P_{29}[(2, 1); Z, \sigma^e, -]$




$P_{29}[(3, 1); Z, \sigma^e, -]$




$P_{29}[(1, 1); Z, \sigma^e, +]$



$P_{29}[(2, 1); Z, \sigma^e, +]$



$P_{29}[(3, 1); Z, \sigma^e, +]$



Summary

- **electron propagations** in the triangular armchair and zigzag graphene quantum dots surrounded by the Neumann and Dirichlet walls determined by the **four Hamiltonians**
- **honeycomb Weyl orbit functions** and the corresponding **discrete Fourier–Weyl transforms**
- **energy spectra** corresponding to the electron armchair and zigzag stationary states exactly calculated
- armchair **Neumann boundary effect**, armchair energy relations transform exactly to the standard graphene tight-binding next-to-nearest neighbour coupling **energy bands**
- zigzag probability **density role switching** between the Dirichlet/Neumann boundary behaviour and the valence/conduction states
- (ideal) zigzag **edge states, repulsive** behaviour of **Neumann walls** predicted for zigzag conduction states
- effects of the three corner positions, zero energy states, higher-degree couplings, energy degeneracy, magnetic field, representations of $SU(3)$



References

- A. V. Rozhkov, F. Nori, *Exact wave functions for an electron on a graphene triangular quantum dot*, Phys. Rev. B **81** (2010) 155401
- J. Akola, H. P. Heiskanen, M. Manninen, *Edge-dependent selection rules in magic triangular graphene flakes*, Phys. Rev. B **77** (2008) 193410
- A. H. Castro Neto, F. Guinea, N. M. R. Peres, K. S. Novoselov, A. K. Geim, *The electronic properties of graphene*, Rev. Mod. Phys. **81** (2009) 109–162
- S. Phark et al., *Direct Observation of Electron Confinement in Epitaxial Graphene Nanoislands*, ACS Nano **5** (2011) 8162–8166
- M. Zarenia, A. Chaves, G. A. Farias, F. M. Peeters, *Energy levels of triangular and hexagonal graphene quantum dots: A comparative study between the tight-binding and Dirac equation approach*, Phys. Rev. B **84** (2011) 245403
- Q. Zhang et al., *Investigation of edge states in artificial graphene nano-flakes*, J. Phys.: Condens. Matter **33** (2021) 225003

