

# Analytic and Algebraic Methods in Physics XXI

*Prague, August 27-30, 2024*

## Some recent applications of the Generalized Uncertainty Principle



**L.M. Nieto**

(collaboration with M. Baradaran and S. Zarrinkamar)

Departamento de Física Teórica, Atómica y Óptica, Univ. Valladolid (Spain)



Financiado por  
la Unión Europea  
NextGenerationEU



Plan de Recuperación,  
Transformación y Resiliencia



- 1 Introduction & Motivation
- 2 Quantum Correction to the Coulomb Potential in GUP Approach
- 3 Dirac equation with generalized gravitational interaction

# 1.– Introduction & Motivation

# 1.– Introduction & Motivation

Due to the importance of a **unified theory of quantum mechanics + gravity**, and the existence of a minimal length (**Planck scale**), we consider

# 1.– Introduction & Motivation

Due to the importance of a **unified theory of quantum mechanics + gravity**, and the existence of a minimal length (**Planck scale**), we consider

- A **modified Schrödinger equation from a generalized uncertainty principle (GUP)**, with a quantum mechanically corrected gravitational interaction. The resulting equation cannot be solved by common exact approaches  $\Rightarrow$  a **Bethe-ansatz** approach.

# 1.– Introduction & Motivation

Due to the importance of a **unified theory of quantum mechanics + gravity**, and the existence of a minimal length (**Planck scale**), we consider

- A **modified Schrödinger equation from a generalized uncertainty principle (GUP)**, with a quantum mechanically corrected gravitational interaction. The resulting equation cannot be solved by common exact approaches  $\Rightarrow$  a **Bethe-ansatz** approach.
  - **Dirac equation with a generalized gravitational interaction** which includes post-Newtonian (relativistic) and quantum corrections to the classical potential.  
The **Bethe-ansatz** approach is also proposed to attack this challenging problem.
-

## 2.– Quantum Correction to the Coulomb Potential in GUP

## 2.– Quantum Correction to the Coulomb Potential in GUP

Several theories provide a unified description of quantum mechanics and gravity: quantum gravity, black hole physics, double special relativity, string theory...



## 2.– Quantum Correction to the Coulomb Potential in GUP

Several theories provide a unified description of quantum mechanics and gravity: quantum gravity, black hole physics, double special relativity, string theory...

The common point in such theories is that they all predict the existence of a minimal length of the Planck scale

$$\ell_p = \sqrt{\hbar G/c^3}.$$

## 2.– Quantum Correction to the Coulomb Potential in GUP

Several theories provide a unified description of quantum mechanics and gravity: quantum gravity, black hole physics, double special relativity, string theory...

The common point in such theories is that they all predict the existence of a minimal length of the Planck scale

$$\ell_p = \sqrt{\hbar G/c^3}.$$

This minimal length is equivalent to a **generalization of the uncertainty principle (GUP)** that affects the entire physical system.

## 2.– Quantum Correction to the Coulomb Potential in GUP

Several theories provide a unified description of quantum mechanics and gravity: quantum gravity, black hole physics, double special relativity, string theory...

The common point in such theories is that they all predict the existence of a minimal length of the Planck scale

$$\ell_p = \sqrt{\hbar G/c^3}.$$

This minimal length is equivalent to a **generalization of the uncertainty principle (GUP)** that affects the entire physical system.

From the mathematical point of view, in a GUP formalism the **Schrödinger equation is of higher order** four, six or more, depending on the choice of the GUP and the operators involved.

## 2.– Quantum Correction to the Coulomb Potential in GUP

Several theories provide a unified description of quantum mechanics and gravity: quantum gravity, black hole physics, double special relativity, string theory...

The common point in such theories is that they all predict the existence of a minimal length of the Planck scale

$$\ell_p = \sqrt{\hbar G/c^3}.$$

This minimal length is equivalent to a **generalization of the uncertainty principle (GUP)** that affects the entire physical system.

From the mathematical point of view, in a GUP formalism the **Schrödinger equation is of higher order** four, six or more, depending on the choice of the GUP and the operators involved.

Obviously, those are not well studied problems in mathematical physics, where the differential equations that appear, whether relativistic or non-relativistic, are of the first or second order.

## 2.– Quantum Correction to the Coulomb Potential in GUP

On the other hand, the **Coulomb/Kepler potential** is obviously the most physical interaction in common with gravity and quantum mechanics.

## 2.– Quantum Correction to the Coulomb Potential in GUP

On the other hand, the **Coulomb/Kepler potential** is obviously the most physical interaction in common with gravity and quantum mechanics.

The quantum Coulomb interaction remains an **attractive and challenging field of study** and very recent articles continue to discuss related properties.

## 2.– Quantum Correction to the Coulomb Potential in GUP

On the other hand, the **Coulomb/Kepler potential** is obviously the most physical interaction in common with gravity and quantum mechanics.

The quantum Coulomb interaction remains an **attractive and challenging field of study** and very recent articles continue to discuss related properties.

It has been shown that the quantum correction of Coulomb potential will contain **inverse quadratic, cubic and quartic terms**.

## 2.– Quantum Correction to the Coulomb Potential in GUP

On the other hand, the **Coulomb/Kepler potential** is obviously the most physical interaction in common with gravity and quantum mechanics.

The quantum Coulomb interaction remains an **attractive and challenging field of study** and very recent articles continue to discuss related properties.

It has been shown that the quantum correction of Coulomb potential will contain **inverse quadratic, cubic and quartic terms**.

This is of particular interest to us: what we are going to do next is to analyze this type of corrected Coulomb potential that includes negative powers up to the fourth order, which we will call **Coulomb–4** potential, within the GUP formalism.



## 2.– Quantum Correction to the Coulomb Potential in GUP

On the other hand, the **Coulomb/Kepler potential** is obviously the most physical interaction in common with gravity and quantum mechanics.

The quantum Coulomb interaction remains an **attractive and challenging field of study** and very recent articles continue to discuss related properties.

It has been shown that the quantum correction of Coulomb potential will contain **inverse quadratic, cubic and quartic terms**.

This is of particular interest to us: what we are going to do next is to analyze this type of corrected Coulomb potential that includes negative powers up to the fourth order, which we will call **Coulomb–4** potential, within the GUP formalism.

**Roadmap:**

## 2.– Quantum Correction to the Coulomb Potential in GUP

On the other hand, the **Coulomb/Kepler potential** is obviously the most physical interaction in common with gravity and quantum mechanics.

The quantum Coulomb interaction remains an **attractive and challenging field of study** and very recent articles continue to discuss related properties.

It has been shown that the quantum correction of Coulomb potential will contain **inverse quadratic, cubic and quartic terms**.

This is of particular interest to us: what we are going to do next is to analyze this type of corrected Coulomb potential that includes negative powers up to the fourth order, which we will call **Coulomb–4** potential, within the GUP formalism.

### **Roadmap:**

(a) **Schrödinger equation modified with GUP** formalism is reviewed compactly;

## 2.– Quantum Correction to the Coulomb Potential in GUP

On the other hand, the **Coulomb/Kepler potential** is obviously the most physical interaction in common with gravity and quantum mechanics.

The quantum Coulomb interaction remains an **attractive and challenging field of study** and very recent articles continue to discuss related properties.

It has been shown that the quantum correction of Coulomb potential will contain **inverse quadratic, cubic and quartic terms**.

This is of particular interest to us: what we are going to do next is to analyze this type of corrected Coulomb potential that includes negative powers up to the fourth order, which we will call **Coulomb–4** potential, within the GUP formalism.

### Roadmap:

- (a) **Schrödinger equation modified with GUP** formalism is reviewed compactly;
- (b) Solutions to the ordinary case (**without GUP**) using the Lie algebraic approach and Heun functions are investigated;

## 2.– Quantum Correction to the Coulomb Potential in GUP

On the other hand, the **Coulomb/Kepler potential** is obviously the most physical interaction in common with gravity and quantum mechanics.

The quantum Coulomb interaction remains an **attractive and challenging field of study** and very recent articles continue to discuss related properties.

It has been shown that the quantum correction of Coulomb potential will contain **inverse quadratic, cubic and quartic terms**.

This is of particular interest to us: what we are going to do next is to analyze this type of corrected Coulomb potential that includes negative powers up to the fourth order, which we will call **Coulomb–4** potential, within the GUP formalism.

### Roadmap:

- (a) **Schrödinger equation modified with GUP** formalism is reviewed compactly;
- (b) Solutions to the ordinary case (**without GUP**) using the Lie algebraic approach and Heun functions are investigated;
- (c) Solution to the **modified GUP problem** are explored;

## 2.– Quantum Correction to the Coulomb Potential in GUP

On the other hand, the **Coulomb/Kepler potential** is obviously the most physical interaction in common with gravity and quantum mechanics.

The quantum Coulomb interaction remains an **attractive and challenging field of study** and very recent articles continue to discuss related properties.

It has been shown that the quantum correction of Coulomb potential will contain **inverse quadratic, cubic and quartic terms**.

This is of particular interest to us: what we are going to do next is to analyze this type of corrected Coulomb potential that includes negative powers up to the fourth order, which we will call **Coulomb–4** potential, within the GUP formalism.

### Roadmap:

- (a) **Schrödinger equation modified with GUP** formalism is reviewed compactly;
- (b) Solutions to the ordinary case (**without GUP**) using the Lie algebraic approach and Heun functions are investigated;
- (c) Solution to the **modified GUP problem** are explored;
- (d) **Ground and first excited states** are explicitly determined.

## 2.– Quantum Correction to the Coulomb Potential in GUP

## 2.– Quantum Correction to the Coulomb Potential in GUP

We start considering a generalized uncertainty principle (**GUP**) of the form

$$[x_G, p_G] = i\hbar \left( 1 + \frac{\beta}{2\hbar^2} p^2 \right), \quad 0 \leq \beta \leq 1,$$

where  $\beta$  is the original minimal length parameter.

## 2.– Quantum Correction to the Coulomb Potential in GUP

We start considering a generalized uncertainty principle (**GUP**) of the form

$$[x_G, p_G] = i\hbar \left( 1 + \frac{\beta}{2\hbar^2} p^2 \right), \quad 0 \leq \beta \leq 1,$$

where  $\beta$  is the original minimal length parameter.

The generalized  $x$  operator is defined as  $x_G = x$ , with  $x$  and  $p$  being the ordinary position and momentum operators, respectively.



## 2.– Quantum Correction to the Coulomb Potential in GUP

We start considering a generalized uncertainty principle (**GUP**) of the form

$$[x_G, p_G] = i\hbar \left( 1 + \frac{\beta}{2\hbar^2} p^2 \right), \quad 0 \leq \beta \leq 1,$$

where  $\beta$  is the original minimal length parameter.

The generalized  $x$  operator is defined as  $x_G = x$ , with  $x$  and  $p$  being the ordinary position and momentum operators, respectively.

In one spatial dimension, and neglecting some high order terms, the above GUP corresponds to the **modified Schrödinger equation**

$$\left( -\frac{d^2}{dx^2} + V_e(x) \right) \psi_n^G(x) = 0,$$

## 2.– Quantum Correction to the Coulomb Potential in GUP

We start considering a generalized uncertainty principle (**GUP**) of the form

$$[x_G, p_G] = i\hbar \left( 1 + \frac{\beta}{2\hbar^2} p^2 \right), \quad 0 \leq \beta \leq 1,$$

where  $\beta$  is the original minimal length parameter.

The generalized  $x$  operator is defined as  $x_G = x$ , with  $x$  and  $p$  being the ordinary position and momentum operators, respectively.

In one spatial dimension, and neglecting some high order terms, the above GUP corresponds to the **modified Schrödinger equation**

$$\left( -\frac{d^2}{dx^2} + V_e(x) \right) \psi_n^G(x) = 0,$$

with an effective potential

$$V_e(x) := \frac{2m}{\hbar^2} (V(x) - E_n^G) + \left( \frac{2m}{\hbar^2} \right)^2 \beta (V(x) - E_n)^2,$$

with  $E_n$  the  $\beta = 0$  eigenvalues of the energy and  $E_n^G$  the eigenenergies for  $\beta \neq 0$ .

## 2.– Quantum Correction to the Coulomb Potential in GUP

## 2.– Quantum Correction to the Coulomb Potential in GUP

We are interested in analyzing a specific problem that could be of great physical interest: the following quantum correction to the Coulomb interaction that includes negative powers up to fourth order, that we will call Coulomb–4 potential:

$$\frac{2m}{\hbar^2} V(x) = \frac{\alpha_1}{x} + \frac{\alpha_2}{x^2} + \frac{\alpha_3}{x^3} + \frac{\alpha_4^2}{x^4}, \quad \alpha_4 > 0, \alpha_1 < 0,$$

## 2.– Quantum Correction to the Coulomb Potential in GUP

We are interested in analyzing a specific problem that could be of great physical interest: the following quantum correction to the Coulomb interaction that includes negative powers up to fourth order, that we will call Coulomb–4 potential:

$$\frac{2m}{\hbar^2} V(x) = \frac{\alpha_1}{x} + \frac{\alpha_2}{x^2} + \frac{\alpha_3}{x^3} + \frac{\alpha_4^2}{x^4}, \quad \alpha_4 > 0, \alpha_1 < 0,$$

which corresponds to the effective potential

$$V_e(x) = \gamma_0 + \frac{\gamma_1}{x} + \frac{\gamma_2}{x^2} + \frac{\gamma_3}{x^3} + \frac{\gamma_4}{x^4} + \frac{\gamma_5}{x^5} + \frac{\gamma_6}{x^6} + \frac{\gamma_7}{x^7} + \frac{\gamma_8}{x^8},$$

## 2.– Quantum Correction to the Coulomb Potential in GUP

We are interested in analyzing a **specific problem** that could be of great physical interest: the following quantum correction to the Coulomb interaction that includes negative powers up to fourth order, **that we will call Coulomb–4** potential:

$$\frac{2m}{\hbar^2} V(x) = \frac{\alpha_1}{x} + \frac{\alpha_2}{x^2} + \frac{\alpha_3}{x^3} + \frac{\alpha_4^2}{x^4}, \quad \alpha_4 > 0, \alpha_1 < 0,$$

which corresponds to the effective potential

$$V_e(x) = \gamma_0 + \frac{\gamma_1}{x} + \frac{\gamma_2}{x^2} + \frac{\gamma_3}{x^3} + \frac{\gamma_4}{x^4} + \frac{\gamma_5}{x^5} + \frac{\gamma_6}{x^6} + \frac{\gamma_7}{x^7} + \frac{\gamma_8}{x^8},$$

in which the  $\gamma_i$ ,  $i = 0, 1, \dots, 8$  are as follows

$$\begin{aligned} \gamma_0 &= \beta \epsilon_n^2 - \epsilon_n^G, & \gamma_1 &= \alpha_1(1 - 2\beta \epsilon_n), \\ \gamma_2 &= \alpha_2 + \beta(\alpha_1^2 - 2\alpha_2 \epsilon_n), & \gamma_3 &= \alpha_3 + 2\beta(\alpha_1 \alpha_2 - \alpha_3 \epsilon_n), \\ \gamma_4 &= \alpha_4^2 + \beta(2\alpha_1 \alpha_3 + \alpha_2^2 - 2\alpha_4^2 \epsilon_n), & \gamma_5 &= 2\beta(\alpha_1 \alpha_4^2 + \alpha_2 \alpha_3), \\ \gamma_6 &= \beta(2\alpha_2 \alpha_4^2 + \alpha_3^2), & \gamma_7 &= 2\beta \alpha_3 \alpha_4^2, & \gamma_8 &= \beta \alpha_4^4, \end{aligned}$$

where we have introduced the notation

$$\epsilon_n = \frac{2m}{\hbar^2} E_n, \quad \epsilon_n^G = \frac{2m}{\hbar^2} E_n^G.$$

## 2.– Quantum Correction to the Coulomb Potential in GUP

This problem only makes sense in the half-line, that is,  $x$  is always positive and at the origin it is assumed that there is an impenetrable infinite wall that prevents the passage of the particle in the other direction.

## 2.– Quantum Correction to the Coulomb Potential in GUP

This problem only makes sense in the half-line, that is,  $x$  is always positive and at the origin it is assumed that there is an impenetrable infinite wall that prevents the passage of the particle in the other direction.

Before continuing, it is worth briefly commenting on the choice Coulomb–4 interaction:



## 2.– Quantum Correction to the Coulomb Potential in GUP

This problem only makes sense in the half-line, that is,  $x$  is always positive and at the origin it is assumed that there is an impenetrable infinite wall that prevents the passage of the particle in the other direction.

Before continuing, it is worth briefly commenting on the choice Coulomb–4 interaction:

- The first term is the ordinary Coulomb or gravitational potential:  $\frac{\alpha_1}{x}$

## 2.– Quantum Correction to the Coulomb Potential in GUP

This problem only makes sense in the half-line, that is,  $x$  is always positive and at the origin it is assumed that there is an impenetrable infinite wall that prevents the passage of the particle in the other direction.

Before continuing, it is worth briefly commenting on the choice Coulomb–4 interaction:

- The first term is the ordinary Coulomb or gravitational potential:  $\frac{\alpha_1}{x}$
- The second term,  $\frac{\alpha_2}{x^2}$ , is usually considered to be the relativistic correction to the gravitational problem and includes the speed of light,  $c$ .

## 2.– Quantum Correction to the Coulomb Potential in GUP

This problem only makes sense in the half-line, that is,  $x$  is always positive and at the origin it is assumed that there is an impenetrable infinite wall that prevents the passage of the particle in the other direction.

Before continuing, it is worth briefly commenting on the choice Coulomb–4 interaction:

- The first term is the ordinary Coulomb or gravitational potential:  $\frac{\alpha_1}{x}$
- The second term,  $\frac{\alpha_2}{x^2}$ , is usually considered to be the relativistic correction to the gravitational problem and includes the speed of light,  $c$ .
- Regarding the quantum correction, there is not a quite unified approach, but almost all existing ones include the inverse cubic term  $\frac{\alpha_3}{x^3}$  as the necessary correction to the gravitational potential.

## 2.– Quantum Correction to the Coulomb Potential in GUP

This problem only makes sense in the half-line, that is,  $x$  is always positive and at the origin it is assumed that there is an impenetrable infinite wall that prevents the passage of the particle in the other direction.

Before continuing, it is worth briefly commenting on the choice Coulomb–4 interaction:

- The first term is the ordinary Coulomb or gravitational potential:  $\frac{\alpha_1}{x}$
- The second term,  $\frac{\alpha_2}{x^2}$ , is usually considered to be the relativistic correction to the gravitational problem and includes the speed of light,  $c$ .
- Regarding the quantum correction, there is not a quite unified approach, but almost all existing ones include the inverse cubic term  $\frac{\alpha_3}{x^3}$  as the necessary correction to the gravitational potential.

It should be noted that both the sign and the value of the parameters are quite different in several articles, which motivates further studies.

## 2.– Quantum Correction to the Coulomb Potential in GUP

This problem only makes sense in the half-line, that is,  $x$  is always positive and at the origin it is assumed that there is an impenetrable infinite wall that prevents the passage of the particle in the other direction.

Before continuing, it is worth briefly commenting on the choice Coulomb–4 interaction:

- The first term is the ordinary Coulomb or gravitational potential:  $\frac{\alpha_1}{x}$
- The second term,  $\frac{\alpha_2}{x^2}$ , is usually considered to be the relativistic correction to the gravitational problem and includes the speed of light,  $c$ .
- Regarding the quantum correction, there is not a quite unified approach, but almost all existing ones include the inverse cubic term  $\frac{\alpha_3}{x^3}$  as the necessary correction to the gravitational potential.

It should be noted that both the sign and the value of the parameters are quite different in several articles, which motivates further studies.

Let us start the analysis of the ordinary case ( $\beta = 0$ ), as the solutions of the model with GUP depend on them.

## 2.– Quantum Correction to the Coulomb Potential in GUP

**The ordinary case  $\beta = 0$  (not GUP)** has already been studied via the ansatz method. Here, we will obtain the general solutions of the model using the Lie-algebraic method within the framework of quasi-exact solvability.

## 2.– Quantum Correction to the Coulomb Potential in GUP

**The ordinary case  $\beta = 0$  (not GUP)** has already been studied via the ansatz method. Here, we will obtain the general solutions of the model using the Lie-algebraic method within the framework of quasi-exact solvability.

The Schrödinger equation with Coulomb–4 potential appears in the form

$$\left( -\frac{d^2}{dx^2} + \frac{\alpha_1}{x} + \frac{\alpha_2}{x^2} + \frac{\alpha_3}{x^3} + \frac{\alpha_4^2}{x^4} - \epsilon_n \right) \psi_n(x) = 0.$$

## 2.– Quantum Correction to the Coulomb Potential in GUP

**The ordinary case  $\beta = 0$  (not GUP)** has already been studied via the ansatz method. Here, we will obtain the general solutions of the model using the Lie-algebraic method within the framework of quasi-exact solvability.

The Schrödinger equation with Coulomb-4 potential appears in the form

$$\left( -\frac{d^2}{dx^2} + \frac{\alpha_1}{x} + \frac{\alpha_2}{x^2} + \frac{\alpha_3}{x^3} + \frac{\alpha_4^2}{x^4} - \epsilon_n \right) \psi_n(x) = 0.$$

Due to the asymptotic behavior of the wave function  $\psi_n(x)$ , we use the ansatz

$$\psi_n(x) = x^\delta \exp \left[ - \left( x \sqrt{-\epsilon_n} + \frac{\alpha_4}{x} \right) \right] \varphi_n(x), \quad \delta = 1 + \frac{\alpha_3}{2\alpha_4} > 0,$$



## 2.– Quantum Correction to the Coulomb Potential in GUP

**The ordinary case  $\beta = 0$  (not GUP)** has already been studied via the ansatz method. Here, we will obtain the general solutions of the model using the Lie-algebraic method within the framework of quasi-exact solvability.

The Schrödinger equation with Coulomb–4 potential appears in the form

$$\left( -\frac{d^2}{dx^2} + \frac{\alpha_1}{x} + \frac{\alpha_2}{x^2} + \frac{\alpha_3}{x^3} + \frac{\alpha_4^2}{x^4} - \epsilon_n \right) \psi_n(x) = 0.$$

Due to the asymptotic behavior of the wave function  $\psi_n(x)$ , we use the ansatz

$$\psi_n(x) = x^\delta \exp \left[ - \left( x \sqrt{-\epsilon_n} + \frac{\alpha_4}{x} \right) \right] \varphi_n(x), \quad \delta = 1 + \frac{\alpha_3}{2\alpha_4} > 0,$$

which transforms Schrödinger equation (9) into the form

$$\left\{ -x^2 \frac{d^2}{dx^2} - 2 \left( \alpha_4 + \delta x - \sqrt{-\epsilon_n} x^2 \right) \frac{d}{dx} + (\lambda_1 x + \lambda_2) \right\} \varphi_n(x) = 0,$$

where

$$\lambda_1 = \alpha_1 + \left( 2 + \frac{\alpha_3}{\alpha_4} \right) \sqrt{-\epsilon_n}, \quad \lambda_2 = \alpha_2 - \frac{\alpha_3^2}{4\alpha_4^2} - \frac{\alpha_3}{2\alpha_4} + 2\alpha_4 \sqrt{-\epsilon_n}.$$

## 2.– Quantum Correction to the Coulomb Potential in GUP

We are going to analyze the problem from two different and complementary points of view: first the [Lie algebraic approach](#) and then we will use the [Heun functions](#).

## 2.– Quantum Correction to the Coulomb Potential in GUP

We are going to analyze the problem from two different and complementary points of view: first the [Lie algebraic approach](#) and then we will use the [Heun functions](#).

**The Lie-algebraic approach:**

## 2.– Quantum Correction to the Coulomb Potential in GUP

We are going to analyze the problem from two different and complementary points of view: first the **Lie algebraic approach** and then we will use the **Heun functions**.

**The Lie-algebraic approach:** Following the standard idea of quasi-exact solvability, we find that if the constraint

$$\lambda_1 = -2n\sqrt{-\epsilon_n}$$

holds, the Schrödinger equation can then be expressed as a quasi-exactly solvable (QES) differential operator in the Lie-algebraic form

$$H_{ges} \varphi_n(x) = 0,$$

## 2.– Quantum Correction to the Coulomb Potential in GUP

We are going to analyze the problem from two different and complementary points of view: first the **Lie algebraic approach** and then we will use the **Heun functions**.

**The Lie-algebraic approach:** Following the standard idea of quasi-exact solvability, we find that if the constraint

$$\lambda_1 = -2n\sqrt{-\epsilon_n}$$

holds, the Schrödinger equation can then be expressed as a quasi-exactly solvable (QES) differential operator in the Lie-algebraic form

$$H_{ges} \varphi_n(x) = 0,$$

with

$$H_{ges} = -\mathcal{J}_n^+ \mathcal{J}_n^- + 2\sqrt{-\epsilon_n} \mathcal{J}_n^+ - 2\alpha_4 \mathcal{J}_n^- - (2\delta + n) \mathcal{J}_n^0 - \frac{n^2}{2} - n\delta + \lambda_2.$$

Here,

$$\mathcal{J}_n^+ = x^2 \frac{d}{dx} - nx, \quad \mathcal{J}_n^0 = x \frac{d}{dx} - \frac{n}{2}, \quad \mathcal{J}_n^- = \frac{d}{dx},$$

are the generators of the  $sl(2)$  Lie algebra satisfying the commutation relations

$$[\mathcal{J}_n^+, \mathcal{J}_n^-] = -2\mathcal{J}_n^0, \quad [\mathcal{J}_n^\pm, \mathcal{J}_n^0] = \mp \mathcal{J}_n^\pm.$$

## 2.– Quantum Correction to the Coulomb Potential in GUP

These operators leave invariant the  $(n+1)$ -dimensional linear space of polynomials

$$\varphi_n(x) = \sum_{k=0}^n c_k x^k,$$

where the coefficients  $c_k$  satisfy the three-term recursion relation ( $c_{-1} = c_{n+1} = 0$ )

$$c_{k+1} = \frac{(\lambda_2 - 2k\delta - k(k-1))c_k - 4\sqrt{-\epsilon_n}c_{k-1}}{2(k+1)\alpha_4}, \quad k = 0, 1, \dots, n,$$

assuming  $c_{-1} = 0$  and  $c_{n+1} = 0$ .

## 2.– Quantum Correction to the Coulomb Potential in GUP

These operators leave invariant the  $(n+1)$ -dimensional linear space of polynomials

$$\varphi_n(x) = \sum_{k=0}^n c_k x^k,$$

where the coefficients  $c_k$  satisfy the three-term recursion relation ( $c_{-1} = c_{n+1} = 0$ )

$$c_{k+1} = \frac{(\lambda_2 - 2k\delta - k(k-1))c_k - 4\sqrt{-\epsilon_n}c_{k-1}}{2(k+1)\alpha_4}, \quad k = 0, 1, \dots, n,$$

assuming  $c_{-1} = 0$  and  $c_{n+1} = 0$ . Equivalently, the recursion relation (11) can be rewritten as a tridiagonal matrix equation the nontrivial solutions of which are

$$\begin{vmatrix} \lambda_2 & -2\alpha_4 & & & \\ -2n\sqrt{-\epsilon_n} & (\lambda_2 - 2\delta) & -4\alpha_4 & & \\ & -2(n-1)\sqrt{-\epsilon_n} & \ddots & \ddots & \\ & & \ddots & \ddots & -2n\alpha_4 \\ & & & -2\sqrt{-\epsilon_n} & (n - n^2 + \lambda_2 - 2n\delta) \end{vmatrix} = 0.$$

## 2.– Quantum Correction to the Coulomb Potential in GUP

Note that the last condition imposes severe restrictions on the potential parameters  $\alpha_i$ ,  $i = 1, 2, 3, 4$ .



## 2.– Quantum Correction to the Coulomb Potential in GUP

Note that the last condition imposes severe restrictions on the potential parameters  $\alpha_i$ ,  $i = 1, 2, 3, 4$ .

On the other hand, from the expressions of  $\lambda_1$  and  $\lambda_2$  the following expression of the energy in closed form can be obtained

$$\epsilon_n = -\frac{\alpha_1^2 \alpha_4^2}{(\alpha_3 + 2(n+1)\alpha_4)^2},$$

provided  $\alpha_1 < 0$ .

## 2.– Quantum Correction to the Coulomb Potential in GUP

Note that the last condition imposes severe restrictions on the potential parameters  $\alpha_i$ ,  $i = 1, 2, 3, 4$ .

On the other hand, from the expressions of  $\lambda_1$  and  $\lambda_2$  the following expression of the energy in closed form can be obtained

$$\epsilon_n = -\frac{\alpha_1^2 \alpha_4^2}{(\alpha_3 + 2(n+1)\alpha_4)^2},$$

provided  $\alpha_1 < 0$ .

For the sake of clarity, next we will find the explicit solutions of the ground state and the first excited state.


## 2.– Quantum Correction to the Coulomb Potential in GUP

### Ground state and the associated wave function

$$\epsilon_0 = -\frac{\alpha_1^2 \alpha_4^2}{(\alpha_3 + 2\alpha_4)^2}, \quad \psi_0(x) = c_0 x^{1+\alpha_3/(2\alpha_4)} \exp \left[ - \left( x\sqrt{-\epsilon_0} + \frac{\alpha_4}{x} \right) \right],$$

where the restriction on the parameters of the potential is determined by

$$4\alpha_2 \alpha_4^2 - \alpha_3^2 - 2\alpha_3 \alpha_4 + 8\alpha_4^3 \sqrt{-\epsilon_0} = 0.$$



$$\alpha_1 = -1/10$$


## 2.– Quantum Correction to the Coulomb Potential in GUP

To better capture the meaning of the analytical results found above, next Figure represents:

## 2.– Quantum Correction to the Coulomb Potential in GUP

To better capture the meaning of the analytical results found above, next Figure represents:


On the left, the wave function of the ground state, together with the corresponding energy,



## 2.– Quantum Correction to the Coulomb Potential in GUP

To better capture the meaning of the analytical results found above, next Figure represents:

On the left, the wave function of the ground state, together with the corresponding energy,




On the right, the form of the potential.


## 2.– Quantum Correction to the Coulomb Potential in GUP

To better capture the meaning of the analytical results found above, next Figure represents:

On the left, the wave function of the ground state, together with the corresponding energy,



On the right, the form of the potential.



## 2.– Quantum Correction to the Coulomb Potential in GUP

### First excited state ( $n = 1$ ) and the associated wave function

The energy and the corresponding wave function are given by

$$\epsilon_1 = -\frac{\alpha_1^2 \alpha_4^2}{(\alpha_3 + 4\alpha_4)^2}, \quad \psi_1(x) = (c_0 + c_1 x) x^{1+\alpha_3/(2\alpha_4)} \exp \left[ -\left( x\sqrt{-\epsilon_1} + \frac{\alpha_4}{x} \right) \right].$$



## 2.– Quantum Correction to the Coulomb Potential in GUP

### First excited state ( $n = 1$ ) and the associated wave function

The energy and the corresponding wave function are given by

$$\epsilon_1 = -\frac{\alpha_1^2 \alpha_4^2}{(\alpha_3 + 4\alpha_4)^2}, \quad \psi_1(x) = (c_0 + c_1 x) x^{1+\alpha_3/(2\alpha_4)} \exp \left[ -\left( x\sqrt{-\epsilon_1} + \frac{\alpha_4}{x} \right) \right].$$

The restriction on the parameters of the potential is given by

$$64\alpha_1^2\alpha_4^8 + 16\alpha_4^4(\alpha_3 + 4\alpha_4)(\alpha_3^2 + 4\alpha_3\alpha_4 + 8\alpha_4^2 - 4\alpha_2\alpha_4^2)\alpha_1 \\ + (\alpha_3 + 4\alpha_4)^2(\alpha_3^2 + 2\alpha_3\alpha_4 - 4\alpha_2\alpha_4^2)(\alpha_3^2 + 6\alpha_3\alpha_4 + 8\alpha_4^2 - 4\alpha_2\alpha_4^2) = 0.$$

## 2.– Quantum Correction to the Coulomb Potential in GUP


### First excited state ( $n = 1$ ) and the associated wave function

The energy and the corresponding wave function are given by

$$\epsilon_1 = -\frac{\alpha_1^2 \alpha_4^2}{(\alpha_3 + 4\alpha_4)^2}, \quad \psi_1(x) = (c_0 + c_1 x) x^{1+\alpha_3/(2\alpha_4)} \exp \left[ -\left( x\sqrt{-\epsilon_1} + \frac{\alpha_4}{x} \right) \right].$$

The restriction on the parameters of the potential is given by

$$64\alpha_1^2 \alpha_4^8 + 16\alpha_4^4 (\alpha_3 + 4\alpha_4) (\alpha_3^2 + 4\alpha_3 \alpha_4 + 8\alpha_4^2 - 4\alpha_2 \alpha_4^2) \alpha_1 \\ + (\alpha_3 + 4\alpha_4)^2 (\alpha_3^2 + 2\alpha_3 \alpha_4 - 4\alpha_2 \alpha_4^2) (\alpha_3^2 + 6\alpha_3 \alpha_4 + 8\alpha_4^2 - 4\alpha_2 \alpha_4^2) = 0.$$



$$\alpha_1 = -\frac{1}{5}$$


## 2.– Quantum Correction to the Coulomb Potential in GUP

The next Figure represents:

## 2.– Quantum Correction to the Coulomb Potential in GUP

The next Figure represents:


On the left, the wave function of the first excited state, together with the corresponding energy,



## 2.– Quantum Correction to the Coulomb Potential in GUP

The next Figure represents:

On the left, the wave function of the first excited state, together with the corresponding energy,




On the right, the form of the potential.


## 2.– Quantum Correction to the Coulomb Potential in GUP

The next Figure represents:

On the left, the wave function of the first excited state, together with the corresponding energy,



On the right, the form of the potential.



## 2.– Quantum Correction to the Coulomb Potential in GUP

## 2.– Quantum Correction to the Coulomb Potential in GUP

**The ordinary case as a double-confluent Heun equation:**



## 2.– Quantum Correction to the Coulomb Potential in GUP

**The ordinary case as a double-confluent Heun equation:** By changing the independent variable  $y = 2\sqrt{-\epsilon_n} x$ , the Schrödinger differential equation

$$\left\{ -x^2 \frac{d^2}{dx^2} - 2(\alpha_4 + \delta x - \sqrt{-\epsilon_n} x^2) \frac{d}{dx} + (\lambda_1 x + \lambda_2) \right\} \varphi_n(x) = 0,$$

## 2.– Quantum Correction to the Coulomb Potential in GUP

**The ordinary case as a double-confluent Heun equation:** By changing the independent variable  $y = 2\sqrt{-\epsilon_n} x$ , the Schrödinger differential equation

$$\left\{ -x^2 \frac{d^2}{dx^2} - 2(\alpha_4 + \delta x - \sqrt{-\epsilon_n} x^2) \frac{d}{dx} + (\lambda_1 x + \lambda_2) \right\} \varphi_n(x) = 0,$$

is transformed into the double-confluent Heun equation

$$\left\{ y^2 \frac{d^2}{dy^2} + (-y^2 + \rho y + \eta) \frac{d}{dy} - (\omega y + \lambda_2) \right\} \varphi_n(y) = 0,$$

in which we denote  $\rho = 2 + \frac{\alpha_3}{\alpha_4}$ ,  $\eta = 4\alpha_4\sqrt{-\epsilon_n}$ ,  $\omega = 1 + \frac{\alpha_1}{2\sqrt{-\epsilon_n}} + \frac{\alpha_3}{2\alpha_4}$ .

## 2.– Quantum Correction to the Coulomb Potential in GUP

**The ordinary case as a double-confluent Heun equation:** By changing the independent variable  $y = 2\sqrt{-\epsilon_n} x$ , the Schrödinger differential equation

$$\left\{ -x^2 \frac{d^2}{dx^2} - 2(\alpha_4 + \delta x - \sqrt{-\epsilon_n} x^2) \frac{d}{dx} + (\lambda_1 x + \lambda_2) \right\} \varphi_n(x) = 0,$$

is transformed into the double-confluent Heun equation

$$\left\{ y^2 \frac{d^2}{dy^2} + (-y^2 + \rho y + \eta) \frac{d}{dy} - (\omega y + \lambda_2) \right\} \varphi_n(y) = 0,$$

in which we denote  $\rho = 2 + \frac{\alpha_3}{\alpha_4}$ ,  $\eta = 4\alpha_4\sqrt{-\epsilon_n}$ ,  $\omega = 1 + \frac{\alpha_1}{2\sqrt{-\epsilon_n}} + \frac{\alpha_3}{2\alpha_4}$ .

Its regular solutions at origin are given by ( $h_0 = 1$ )

$$\varphi := \varphi_n(y; \rho, \eta, \omega, \lambda_2) = \sum_{n=0}^{\infty} h_n x^n,$$

$$h_{n+2} = \frac{(\lambda_2 - n(n + \rho + 1) - \rho) h_{n+1} + (n + \omega) h_n}{(n + 2)\eta} \quad \text{and} \quad h_1 = \frac{\lambda_2}{\eta} h_0,$$

Consequently,  $\varphi$  can admit polynomial solution of degree  $m$  if  $(m + \omega)$  and  $h_{m+1}$  vanish simultaneously. In this way, general solutions to the problem can be obtained in terms of the associated Heun functions.

## 2.– Quantum Correction to the Coulomb Potential in GUP

## 2.– Quantum Correction to the Coulomb Potential in GUP

The GUP case,  $\beta \neq 0$ .

## 2.– Quantum Correction to the Coulomb Potential in GUP

### The GUP case, $\beta \neq 0$ .

Having obtained the solutions of the Coulomb–4 model without GUP in the previous section, let us now return to the modified Schrödinger equation in a formalism with GUP, that is, to the equations

$$\left[ -\frac{d^2}{dx^2} + V_e(x) \right] \psi_n^G(x) = 0, \quad V_e(x) := \frac{2m}{\hbar^2} (V(x) - E_n^G) + \left( \frac{2m}{\hbar^2} \right)^2 \beta (V(x) - E_n^G)^2,$$

## 2.– Quantum Correction to the Coulomb Potential in GUP

### The GUP case, $\beta \neq 0$ .

Having obtained the solutions of the Coulomb–4 model without GUP in the previous section, let us now return to the modified Schrödinger equation in a formalism with GUP, that is, to the equations

$$\left[ -\frac{d^2}{dx^2} + V_e(x) \right] \psi_n^G(x) = 0, \quad V_e(x) := \frac{2m}{\hbar^2} (V(x) - E_n^G) + \left( \frac{2m}{\hbar^2} \right)^2 \beta (V(x) - E_n)^2,$$

To ensure **proper asymptotic behavior** of the wave function  $\psi_n^G(x)$ , after inspecting the differential equation we propose the following ansatz

$$\psi_n^G(x) = x^f e^{g(x)} \varphi_n^G(x), \quad g(x) = -a x - \frac{b}{x} - \frac{c}{x^2} - \frac{d}{x^3},$$

## 2.– Quantum Correction to the Coulomb Potential in GUP

### The GUP case, $\beta \neq 0$ .

Having obtained the solutions of the Coulomb–4 model without GUP in the previous section, let us now return to the modified Schrödinger equation in a formalism with GUP, that is, to the equations

$$\left[ -\frac{d^2}{dx^2} + V_e(x) \right] \psi_n^G(x) = 0, \quad V_e(x) := \frac{2m}{\hbar^2} (V(x) - E_n^G) + \left( \frac{2m}{\hbar^2} \right)^2 \beta (V(x) - E_n)^2,$$

To ensure **proper asymptotic behavior** of the wave function  $\psi_n^G(x)$ , after inspecting the differential equation we propose the following ansatz

$$\psi_n^G(x) = x^f e^{g(x)} \varphi_n^G(x), \quad g(x) = -a x - \frac{b}{x} - \frac{c}{x^2} - \frac{d}{x^3},$$

where  $\varphi_n^G(x)$  must be a **polynomial** and the parameters;  $a > 0, b, c, d > 0, f > 0$  are still unknown, but must be such that the original function  $\psi_n^G(x)$  is square integrable.



## 2.– Quantum Correction to the Coulomb Potential in GUP

### The GUP case, $\beta \neq 0$ .

Having obtained the solutions of the Coulomb–4 model without GUP in the previous section, let us now return to the modified Schrödinger equation in a formalism with GUP, that is, to the equations

$$\left[ -\frac{d^2}{dx^2} + V_e(x) \right] \psi_n^G(x) = 0, \quad V_e(x) := \frac{2m}{\hbar^2} (V(x) - E_n^G) + \left( \frac{2m}{\hbar^2} \right)^2 \beta (V(x) - E_n^G)^2,$$

To ensure **proper asymptotic behavior** of the wave function  $\psi_n^G(x)$ , after inspecting the differential equation we propose the following ansatz

$$\psi_n^G(x) = x^f e^{g(x)} \varphi_n^G(x), \quad g(x) = -a x - \frac{b}{x} - \frac{c}{x^2} - \frac{d}{x^3},$$

where  $\varphi_n^G(x)$  must be a **polynomial** and the parameters;  $a > 0, b, c, d > 0, f > 0$  are still unknown, but must be such that the original function  $\psi_n^G(x)$  is square integrable.

It can be seen that the previous parameters are determined by:

## 2.- Quantum Correction to the Coulomb Potential in GUP

$$a = \sqrt{\gamma_0} = \sqrt{\beta \epsilon_n^2 - \epsilon_n^G} > 0,$$

$$b = -\frac{\gamma_7^2 - 4\gamma_6 \gamma_8}{8(\gamma_8)^{3/2}} = \alpha_2 \sqrt{\beta},$$

$$c = \frac{\gamma_7}{4\sqrt{\gamma_8}} = \frac{1}{2} \alpha_3 \sqrt{\beta},$$

$$d = \frac{\sqrt{\gamma_8}}{3} = \frac{1}{3} \alpha_4^2 \sqrt{\beta},$$

$$f = 2 + \frac{8\gamma_5 \gamma_8^2 - 4\gamma_6 \gamma_7 \gamma_8 + \gamma_7^3}{16(\gamma_8)^{5/2}} = 2 + \alpha_1 \sqrt{\beta} > 0.$$

## 2.– Quantum Correction to the Coulomb Potential in GUP

$$\begin{aligned}a &= \sqrt{\gamma_0} = \sqrt{\beta \epsilon_n^2 - \epsilon_n^G} > 0, & b &= -\frac{\gamma_7^2 - 4\gamma_6 \gamma_8}{8(\gamma_8)^{3/2}} = \alpha_2 \sqrt{\beta}, \\c &= \frac{\gamma_7}{4\sqrt{\gamma_8}} = \frac{1}{2} \alpha_3 \sqrt{\beta}, & d &= \frac{\sqrt{\gamma_8}}{3} = \frac{1}{3} \alpha_4^2 \sqrt{\beta}, \\f &= 2 + \frac{8\gamma_5 \gamma_8^2 - 4\gamma_6 \gamma_7 \gamma_8 + \gamma_7^3}{16(\gamma_8)^{5/2}} = 2 + \alpha_1 \sqrt{\beta} > 0.\end{aligned}$$

Then, the differential equation for  $\varphi_n^G(x)$  simplifies to

$$\left\{ P_4(x) \frac{d^2}{dx^2} + Q_4(x) \frac{d}{dx} + W_3(x) \right\} \varphi_n^G(x) = 0,$$

## 2.– Quantum Correction to the Coulomb Potential in GUP

$$\begin{aligned}a &= \sqrt{\gamma_0} = \sqrt{\beta \epsilon_n^2 - \epsilon_n^G} > 0, & b &= -\frac{\gamma_7^2 - 4\gamma_6 \gamma_8}{8(\gamma_8)^{3/2}} = \alpha_2 \sqrt{\beta}, \\c &= \frac{\gamma_7}{4\sqrt{\gamma_8}} = \frac{1}{2} \alpha_3 \sqrt{\beta}, & d &= \frac{\sqrt{\gamma_8}}{3} = \frac{1}{3} \alpha_4^2 \sqrt{\beta}, \\f &= 2 + \frac{8\gamma_5 \gamma_8^2 - 4\gamma_6 \gamma_7 \gamma_8 + \gamma_7^3}{16(\gamma_8)^{5/2}} = 2 + \alpha_1 \sqrt{\beta} > 0.\end{aligned}$$

Then, the differential equation for  $\varphi_n^G(x)$  simplifies to

$$\left\{ P_4(x) \frac{d^2}{dx^2} + Q_4(x) \frac{d}{dx} + W_3(x) \right\} \varphi_n^G(x) = 0,$$

where

$$\begin{aligned}P_4(x) &= x^4, \\Q_4(x) &= 6d + 4cx + 2bx^2 + 2fx^3 - 2ax^4, \\W_3(x) &= (b^2 - 6ad + 4cf - 6c - \gamma_4) + (2(bf - b - 2ac) - \gamma_3)x \\&\quad + (f(f - 1) - 2ab - \gamma_2)x^2 - (\gamma_1 + 2af)x^3.\end{aligned}$$

## 2.– Quantum Correction to the Coulomb Potential in GUP

To find the solution  $\varphi_n^G(x)$  we use the general **Bethe ansatz method**, looking for polynomial solutions of the form

$$\varphi_n^G(x) = \begin{cases} 1, & n = 0, \\ \prod_{i=1}^n (x - x_i), & n \in \mathbb{N}, \end{cases}$$

where  $x_i$  are distinct roots to be determined.

## 2.– Quantum Correction to the Coulomb Potential in GUP

To find the solution  $\varphi_n^G(x)$  we use the general **Bethe ansatz method**, looking for polynomial solutions of the form

$$\varphi_n^G(x) = \begin{cases} 1, & n = 0, \\ \prod_{i=1}^n (x - x_i), & n \in \mathbb{N}, \end{cases}$$

where  $x_i$  are distinct roots to be determined.

After some calculations we get the energy relation: for a given  $n$ , the energy  $\epsilon_n^G$  is

$$\epsilon_n^G(\epsilon_n; \alpha_1, \beta) = -\frac{\alpha_1^2}{4} \left( \frac{2\beta \epsilon_n - 1}{\alpha_1 \sqrt{\beta} + n + 2} \right)^2 + \beta \epsilon_n^2.$$

The denominator of the fraction,  $\alpha_1 \sqrt{\beta} + n + 2$ , is always positive.

## 2.– Quantum Correction to the Coulomb Potential in GUP

To find the solution  $\varphi_n^G(x)$  we use the general **Bethe ansatz method**, looking for polynomial solutions of the form

$$\varphi_n^G(x) = \begin{cases} 1, & n = 0, \\ \prod_{i=1}^n (x - x_i), & n \in \mathbb{N}, \end{cases}$$

where  $x_i$  are distinct roots to be determined.

After some calculations we get the energy relation: for a given  $n$ , the energy  $\epsilon_n^G$  is

$$\epsilon_n^G(\epsilon_n; \alpha_1, \beta) = -\frac{\alpha_1^2}{4} \left( \frac{2\beta \epsilon_n - 1}{\alpha_1 \sqrt{\beta} + n + 2} \right)^2 + \beta \epsilon_n^2.$$

The denominator of the fraction,  $\alpha_1 \sqrt{\beta} + n + 2$ , is always positive.

As in the ordinary case (without a GUP,  $\beta = 0$ ), we will now look for explicit solutions for the ground state and the first excited state.

## 2.– Quantum Correction to the Coulomb Potential in GUP

### Ground state and the associated wave function

For  $n = 0$ , the energy of ground state,  $\epsilon_0^G$ , is determined in closed form:

$$\epsilon_0^G(\epsilon_0; \alpha_1, \beta) = -\frac{\alpha_1^2}{4} \left( \frac{2\beta \epsilon_0 - 1}{\alpha_1 \sqrt{\beta} + 2} \right)^2 + \beta \epsilon_0^2.$$



## 2.– Quantum Correction to the Coulomb Potential in GUP

### Ground state and the associated wave function

For  $n = 0$ , the energy of ground state,  $\epsilon_0^G$ , is determined in closed form:

$$\epsilon_0^G(\epsilon_0; \alpha_1, \beta) = -\frac{\alpha_1^2}{4} \left( \frac{2\beta \epsilon_0 - 1}{\alpha_1 \sqrt{\beta} + 2} \right)^2 + \beta \epsilon_0^2.$$

The explicit form of the associated wave function is given by

$$\psi_0^G(x) = C_0 x^{2+\alpha_1\sqrt{\beta}} \exp \left[ -x \sqrt{\beta \epsilon_0^2 - \epsilon_0^G} - \frac{\sqrt{\beta}}{2} \left( \frac{2\alpha_4^2}{3x^3} + \frac{\alpha_3}{x^2} + \frac{2\alpha_2}{x} \right) \right],$$

where  $C_0$  is the normalization constant and the parameters of the potential satisfy some constraints.

## 2.– Quantum Correction to the Coulomb Potential in GUP

### First excited state and the associated wave function

For  $n = 1$ , the first excited state energy,  $\epsilon_1^G$ , is given by

$$\epsilon_1^G(\epsilon_1; \alpha_1, \beta) = -\frac{\alpha_1^2}{4} \left( \frac{2\beta \epsilon_1 - 1}{\alpha_1 \sqrt{\beta} + 3} \right)^2 + \beta \epsilon_1^2.$$

## 2.– Quantum Correction to the Coulomb Potential in GUP

### First excited state and the associated wave function

For  $n = 1$ , the first excited state energy,  $\epsilon_1^G$ , is given by

$$\epsilon_1^G(\epsilon_1; \alpha_1, \beta) = -\frac{\alpha_1^2}{4} \left( \frac{2\beta \epsilon_1 - 1}{\alpha_1 \sqrt{\beta} + 3} \right)^2 + \beta \epsilon_1^2.$$

The wave function is explicitly given by

$$\psi_1^G(x) = C_1 (x - x_1) x^{2+\alpha_1 \sqrt{\beta}} \exp \left[ -x \sqrt{\beta \epsilon_1^2 - \epsilon_1^G} - \frac{\sqrt{\beta}}{2} \left( \frac{2\alpha_4^2}{3x^3} + \frac{\alpha_3}{x^2} + \frac{2\alpha_2}{x} \right) \right],$$

where  $C_1$  is the normalization constant and the parameters of the potential satisfy some constraints.

### 3.– Dirac eqn. with generalized gravitational interaction

### 3.– Dirac eqn. with generalized gravitational interaction

In the last part of the talk, the **Dirac equation** is considered with the recently proposed **generalized gravitational interaction** (Kepler or Coulomb), which includes post-Newtonian (relativistic) and quantum corrections to the classical potential.

### 3.– Dirac eqn. with generalized gravitational interaction

In the last part of the talk, the **Dirac equation** is considered with the recently proposed **generalized gravitational interaction** (Kepler or Coulomb), which includes post-Newtonian (relativistic) and quantum corrections to the classical potential.

The general idea in choosing the metric is that the spacetime contributions are contained in an external potential or in an electromagnetic potential which can be considered as a good basis for future studies on space quantum communication.

### 3.– Dirac eqn. with generalized gravitational interaction

In the last part of the talk, the **Dirac equation** is considered with the recently proposed **generalized gravitational interaction** (Kepler or Coulomb), which includes post-Newtonian (relativistic) and quantum corrections to the classical potential.

The general idea in choosing the metric is that the spacetime contributions are contained in an external potential or in an electromagnetic potential which can be considered as a good basis for future studies on space quantum communication.

We also discuss several known generalizations of the Coulomb potential within this formulation in terms of certain Heun functions.

### 3.– Dirac eqn. with generalized gravitational interaction

In the last part of the talk, the **Dirac equation** is considered with the recently proposed **generalized gravitational interaction** (Kepler or Coulomb), which includes post-Newtonian (relativistic) and quantum corrections to the classical potential.

The general idea in choosing the metric is that the spacetime contributions are contained in an external potential or in an electromagnetic potential which can be considered as a good basis for future studies on space quantum communication.

We also discuss several known generalizations of the Coulomb potential within this formulation in terms of certain Heun functions.

#### **Roadmap:**



### 3.– Dirac eqn. with generalized gravitational interaction

In the last part of the talk, the **Dirac equation** is considered with the recently proposed **generalized gravitational interaction** (Kepler or Coulomb), which includes post-Newtonian (relativistic) and quantum corrections to the classical potential.

The general idea in choosing the metric is that the spacetime contributions are contained in an external potential or in an electromagnetic potential which can be considered as a good basis for future studies on space quantum communication.

We also discuss several known generalizations of the Coulomb potential within this formulation in terms of certain Heun functions.

#### **Roadmap:**

(A) **A review** of the essential formulae of the Dirac equation in the desired metric.

### 3.– Dirac eqn. with generalized gravitational interaction

In the last part of the talk, the **Dirac equation** is considered with the recently proposed **generalized gravitational interaction** (Kepler or Coulomb), which includes post-Newtonian (relativistic) and quantum corrections to the classical potential.

The general idea in choosing the metric is that the spacetime contributions are contained in an external potential or in an electromagnetic potential which can be considered as a good basis for future studies on space quantum communication.

We also discuss several known generalizations of the Coulomb potential within this formulation in terms of certain Heun functions.

#### **Roadmap:**

- (A) **A review** of the essential formulae of the Dirac equation in the desired metric.
- (B) Using the **Bethe-ansatz approach**, we report the general solution for arbitrary  $n$ , in particular solutions for the ground and the first excited states.

## 3.– Dirac eqn. with generalized gravitational interaction

### (A) Curved spacetime Dirac equation with a generalized metric

### 3.– Dirac eqn. with generalized gravitational interaction

#### (A) Curved spacetime Dirac equation with a generalized metric

The Dirac equation  $(i\gamma^\mu \nabla_\mu - mc) \Psi = 0$  will be considered for the metric

$$g_{\mu\nu} = \text{diag} \left( e^{2f(r)}, -e^{2g(r)}, -r^2, -r^2 \sin^2 \theta \right),$$

### 3.– Dirac eqn. with generalized gravitational interaction

#### (A) Curved spacetime Dirac equation with a generalized metric

The Dirac equation  $(i\gamma^\mu \nabla_\mu - mc)\Psi = 0$  will be considered for the metric

$$g_{\mu\nu} = \text{diag} \left( e^{2f(r)}, -e^{2g(r)}, -r^2, -r^2 \sin^2 \theta \right),$$

where  $f(r)$  and  $g(r)$  are arbitrary functions of the radial coordinate, being the angular parts the same as in (3+1)–Minkowski spacetime.

### 3.– Dirac eqn. with generalized gravitational interaction

#### (A) Curved spacetime Dirac equation with a generalized metric

The Dirac equation  $(i\gamma^\mu \nabla_\mu - mc) \Psi = 0$  will be considered for the metric

$$g_{\mu\nu} = \text{diag} \left( e^{2f(r)}, -e^{2g(r)}, -r^2, -r^2 \sin^2 \theta \right),$$

where  $f(r)$  and  $g(r)$  are arbitrary functions of the radial coordinate, being the angular parts the same as in (3+1)–Minkowski spacetime.

In curved spacetime  $\nabla_\mu = \partial_\mu + iA_\mu/c + \Omega_\mu$ , with  $\partial_\mu$  the covariant derivative on flat spacetime and  $\Omega_\mu$  the spin connection. Choosing  $A_\mu = (V(r), cA_r(r), 0, 0)$ , the spinor wave function is

$$\Psi_c(r, \theta, \phi) = N \begin{pmatrix} R_1(r) \mathcal{Y}_{j+1/2}^{|m|j}(\theta, \phi) \\ iR_2(r) \mathcal{Y}_{j-1/2}^{|m|j}(\theta, \phi) \end{pmatrix}.$$

### 3.– Dirac eqn. with generalized gravitational interaction

#### (A) Curved spacetime Dirac equation with a generalized metric

The Dirac equation  $(i\gamma^\mu \nabla_\mu - mc) \Psi = 0$  will be considered for the metric

$$g_{\mu\nu} = \text{diag} \left( e^{2f(r)}, -e^{2g(r)}, -r^2, -r^2 \sin^2 \theta \right),$$

where  $f(r)$  and  $g(r)$  are arbitrary functions of the radial coordinate, being the angular parts the same as in (3+1)–Minkowski spacetime.

In curved spacetime  $\nabla_\mu = \partial_\mu + iA_\mu/c + \Omega_\mu$ , with  $\partial_\mu$  the covariant derivative on flat spacetime and  $\Omega_\mu$  the spin connection. Choosing  $A_\mu = (V(r), cA_r(r), 0, 0)$ , the spinor wave function is

$$\Psi_c(r, \theta, \phi) = N \begin{pmatrix} R_1(r) \mathcal{Y}_{j+1/2}^{|m|j}(\theta, \phi) \\ iR_2(r) \mathcal{Y}_{j-1/2}^{|m|j}(\theta, \phi) \end{pmatrix}.$$

$$\mathcal{Y}_l^{j=l\pm 1/2 m}(\theta, \phi) = \frac{1}{\sqrt{2l+1}} \begin{pmatrix} \pm \sqrt{l \pm m + \frac{1}{2}} Y_l^{m-1/2}(\theta, \phi) \\ \sqrt{l \mp m + \frac{1}{2}} Y_l^{m+1/2}(\theta, \phi) \end{pmatrix}$$

are the **spinor spherical harmonics**, being  $Y_l^m(\theta, \phi)$  the usual spherical harmonics.

### 3.– Dirac eqn. with generalized gravitational interaction

We manipulate Dirac equation  $(i\gamma^\mu \nabla_\mu - mc) \Psi = 0$  with the assumptions:



### 3.– Dirac eqn. with generalized gravitational interaction

We manipulate Dirac equation  $(i\gamma^\mu \nabla_\mu - mc) \Psi = 0$  with the assumptions:

- $m = \hbar = 1$ ,  $c = 1/\alpha$ , with  $\alpha$  being the fine structure constant.
- $f(r) = g(r)$ , with  $e^{f(r)} = 1 + \alpha^2 U(r)$ ,
- $V(r) = a \mathbf{z}(r)$  and  $U(r) = b \mathbf{z}(r)$ ,
- $R_1(r) = \frac{u(r)}{r} e^{-f(r)/2}$  and  $R_2(r) = \frac{v(r)}{r} e^{-f(r)/2}$ ,
- $\begin{pmatrix} \rho_1(r) \\ \rho_2(r) \end{pmatrix} = \mathbf{U}' \begin{pmatrix} u(r) \\ v(r) \end{pmatrix}$ ,  $\mathbf{U}' = \exp(i\eta\sigma_2)$ ,
- $A_r(r) = \frac{\alpha C}{S} \left[ \frac{V(r)}{C} - U(r) \right] - \frac{\lambda}{r} [1 + \alpha^2 U(r)]$ ,  $C = \cos 2\eta$ ,  $S = \sin 2\eta$ ,

### 3.– Dirac eqn. with generalized gravitational interaction

We manipulate Dirac equation  $(i\gamma^\mu \nabla_\mu - mc) \Psi = 0$  with the assumptions:

- $m = \hbar = 1$ ,  $c = 1/\alpha$ , with  $\alpha$  being the fine structure constant.
- $f(r) = g(r)$ , with  $e^{f(r)} = 1 + \alpha^2 U(r)$ ,
- $V(r) = a \mathbf{z}(r)$  and  $U(r) = b \mathbf{z}(r)$ ,
- $R_1(r) = \frac{u(r)}{r} e^{-f(r)/2}$  and  $R_2(r) = \frac{v(r)}{r} e^{-f(r)/2}$ ,
- $\begin{pmatrix} \rho_1(r) \\ \rho_2(r) \end{pmatrix} = \mathbf{U}' \begin{pmatrix} u(r) \\ v(r) \end{pmatrix}$ ,  $\mathbf{U}' = \exp(i\eta\sigma_2)$ ,
- $A_r(r) = \frac{\alpha C}{S} \left[ \frac{V(r)}{C} - U(r) \right] - \frac{\lambda}{r} [1 + \alpha^2 U(r)]$ ,  $C = \cos 2\eta$ ,  $S = \sin 2\eta$ ,

and we obtain the final equation for the component  $\rho_1(r)$

$$\left[ \frac{d^2}{dr^2} + \frac{\alpha}{S} (aC - b) z'(r) - 2(b + \epsilon a) z(r) - \frac{\alpha^2}{S^2} (aC - b)^2 z^2(r) + \frac{\epsilon^2 - 1}{\alpha^2} \right] \rho_1(r) = 0$$

### 3.– Dirac eqn. with generalized gravitational interaction

## 3.– Dirac eqn. with generalized gravitational interaction

### (B) Solutions by the Bethe-ansatz approach

### 3.– Dirac eqn. with generalized gravitational interaction

#### (B) Solutions by the Bethe-ansatz approach

Let us consider the *potential*  $z(r)$  as in the previous Schrödinger-like equation

$$z(r) = \frac{u}{r} + \frac{v}{r^2} + \frac{w}{r^3}, \quad u, v, w < 0,$$

that has been investigated before in the GUP framework.

## 3.– Dirac eqn. with generalized gravitational interaction

### (B) Solutions by the Bethe-ansatz approach

Let us consider the *potential*  $z(r)$  as in the previous Schrödinger-like equation

$$z(r) = \frac{u}{r} + \frac{v}{r^2} + \frac{w}{r^3}, \quad u, v, w < 0,$$

that has been investigated before in the GUP framework. Substituting into the previous Schrödinger-like equation,

$$\left[ \frac{d^2}{dr^2} + \frac{\epsilon_n^2 - 1}{\alpha^2} - \frac{2u(a\epsilon_n + b)}{r} + \frac{1}{S^2} \left( \frac{\Lambda_2}{r^2} + \frac{\Lambda_3}{r^3} + \frac{\Lambda_4}{r^4} \right) - \frac{\alpha^2(b - aC)^2}{S^2} \left( \frac{w^2}{r^6} + \frac{2vw}{r^5} \right) \right] \rho_{1,n}(r) = 0,$$

### 3.– Dirac eqn. with generalized gravitational interaction

#### (B) Solutions by the Bethe-ansatz approach

Let us consider the *potential*  $z(r)$  as in the previous Schrödinger-like equation

$$z(r) = \frac{u}{r} + \frac{v}{r^2} + \frac{w}{r^3}, \quad u, v, w < 0,$$

that has been investigated before in the GUP framework. Substituting into the previous Schrödinger-like equation,

$$\left[ \frac{d^2}{dr^2} + \frac{\epsilon_n^2 - 1}{\alpha^2} - \frac{2u(a\epsilon_n + b)}{r} + \frac{1}{S^2} \left( \frac{\Lambda_2}{r^2} + \frac{\Lambda_3}{r^3} + \frac{\Lambda_4}{r^4} \right) - \frac{\alpha^2(b - aC)^2}{S^2} \left( \frac{w^2}{r^6} + \frac{2vw}{r^5} \right) \right] \rho_{1,n}(r) = 0,$$

where  $\Lambda_2$ ,  $\Lambda_3$  and  $\Lambda_4$  are some functions of  $u, v, w, a, b, \alpha, S, C, \epsilon_n$ . We have added the index  $n$  to  $\epsilon$  and  $\rho_1(r)$  to distinguish some states from others.

### 3.– Dirac eqn. with generalized gravitational interaction

#### (B) Solutions by the Bethe-ansatz approach

Let us consider the *potential*  $z(r)$  as in the previous Schrödinger-like equation

$$z(r) = \frac{u}{r} + \frac{v}{r^2} + \frac{w}{r^3}, \quad u, v, w < 0,$$

that has been investigated before in the GUP framework. Substituting into the previous Schrödinger-like equation,

$$\left[ \frac{d^2}{dr^2} + \frac{\epsilon_n^2 - 1}{\alpha^2} - \frac{2u(a\epsilon_n + b)}{r} + \frac{1}{S^2} \left( \frac{\Lambda_2}{r^2} + \frac{\Lambda_3}{r^3} + \frac{\Lambda_4}{r^4} \right) - \frac{\alpha^2(b - aC)^2}{S^2} \left( \frac{w^2}{r^6} + \frac{2vw}{r^5} \right) \right] \rho_{1,n}(r) = 0,$$

where  $\Lambda_2$ ,  $\Lambda_3$  and  $\Lambda_4$  are some functions of  $u, v, w, a, b, \alpha, S, C, \epsilon_n$ . We have added the index  $n$  to  $\epsilon$  and  $\rho_1(r)$  to distinguish some states from others.

Now, we propose

$$\rho_{1,n}(r) = e^{\Delta(r)} \mathcal{R}_{1,n}(r), \quad \Delta(r) = \delta \ln r + \frac{\gamma}{r^2} + \frac{\beta}{r} + \lambda r,$$

where  $\mathcal{R}_{1,n}(r)$  is a polynomial, and  $\lambda, \beta, \gamma < 0$ ,  $\delta > 0$  are parameters to be find.



### 3.– Dirac eqn. with generalized gravitational interaction

Consequently, the differential equation for  $\mathcal{R}_{1,n}(r)$  simplifies to

$$\left\{ r^3 \frac{d^2}{dr^2} + (2\lambda r^3 + 2\delta r^2 - 2\beta r - 4\gamma) \frac{d}{dr} + \left( \xi_2 r^2 + \frac{\xi_1}{S^2} r + \frac{\xi_0}{S^2} \right) \right\} \mathcal{R}_{1,n}(r) = 0,$$

where  $\xi_0, \xi_1, \xi_2$  are known functions of  $u, v, w, a, b, \alpha, S, C, \epsilon_n$ .

### 3.– Dirac eqn. with generalized gravitational interaction

Consequently, the differential equation for  $\mathcal{R}_{1,n}(r)$  simplifies to

$$\left\{ r^3 \frac{d^2}{dr^2} + (2\lambda r^3 + 2\delta r^2 - 2\beta r - 4\gamma) \frac{d}{dr} + \left( \xi_2 r^2 + \frac{\xi_1}{S^2} r + \frac{\xi_0}{S^2} \right) \right\} \mathcal{R}_{1,n}(r) = 0,$$

where  $\xi_0, \xi_1, \xi_2$  are known functions of  $u, v, w, a, b, \alpha, S, C, \epsilon_n$ .

To solve this equation, we assume  $\mathcal{R}_{1,n}(r)$  to be a polynomial of the form

$$\mathcal{R}_{1,n}(r) = \begin{cases} 1, & n = 0, \\ \prod_{i=1}^n (r - r_i), & n \in \mathbb{N}, \end{cases}$$

where  $r_i$  are distinct roots to be determined. The general solutions are given by

$$\xi_2 + 2n\lambda = 0,$$

$$\xi_1 + 2\lambda \sum_{i=1}^n r_i + n(n-1) + 2n\delta = 0,$$

$$\xi_0 + 2\lambda \sum_{i=1}^n r_i^2 + 2(\delta + n - 1) \sum_{i=1}^n r_i - 2n\beta = 0,$$

the  $r_i$  given by the Bethe ansatz eqns 
$$\sum_{j=1, j \neq i}^n \frac{1}{r_i - r_j} + \frac{\lambda r_i^3 + \delta r_i^2 - \beta r_i - 2\gamma}{r_i^3} = 0.$$

### 3.– Dirac eqn. with generalized gravitational interaction

#### Ground state and the associated wave function

### 3.– Dirac eqn. with generalized gravitational interaction

#### Ground state and the associated wave function

For  $n = 0$  it follows that the ground state energy  $\epsilon_0$  is given by

$$\epsilon_0 = \frac{-4a\alpha^2\mathcal{A}^2bu^2 \pm \sqrt{4\alpha^2\mathcal{A}^2u^2(a^2 - b^2)(\mathcal{B} - 3\mathcal{A})^2 + (\mathcal{B} - 3\mathcal{A})^4}}{4a^2\alpha^2\mathcal{A}^2u^2 + (\mathcal{B} - 3\mathcal{A})^2},$$

with

$$\mathcal{A} := \sqrt{\alpha^2 S^2 (b - aC)^2}, \quad \mathcal{B} := \alpha(b - aC)(2\alpha u(b - aC) - 3S).$$

### 3.– Dirac eqn. with generalized gravitational interaction

#### Ground state and the associated wave function

For  $n = 0$  it follows that the ground state energy  $\epsilon_0$  is given by

$$\epsilon_0 = \frac{-4a\alpha^2 \mathcal{A}^2 b u^2 \pm \sqrt{4\alpha^2 \mathcal{A}^2 u^2 (a^2 - b^2)(\mathcal{B} - 3\mathcal{A})^2 + (\mathcal{B} - 3\mathcal{A})^4}}{4a^2 \alpha^2 \mathcal{A}^2 u^2 + (\mathcal{B} - 3\mathcal{A})^2},$$

with

$$\mathcal{A} := \sqrt{\alpha^2 S^2 (b - aC)^2}, \quad \mathcal{B} := \alpha(b - aC)(2\alpha u(b - aC) - 3S).$$

The associated wave function is  $\rho_{1,0}(r) \propto e^{\Delta(r)}$ , with

$$\Delta(r) = \left( \frac{3}{2} + \frac{\alpha w(b - aC)(2\alpha u(b - aC) - 3S)}{2\sigma S^2} \right) \ln r - \frac{\sigma}{r^2} \left( \frac{rv}{w} + \frac{1}{2} \right) - \sqrt{\frac{1 - \epsilon_0^2}{\alpha^2}} r,$$

The potential parameters  $v$  and  $w$  are given in terms of  $u, a, b, C, S, \alpha, \epsilon_0$ .

### 3.– Dirac eqn. with generalized gravitational interaction

#### First excited state and the associated wave function

### 3.– Dirac eqn. with generalized gravitational interaction

#### First excited state and the associated wave function

For  $n = 1$ , the first excited-state energy,  $\epsilon_1$ , is given by

$$\epsilon_1 = \frac{-4a\alpha^2\mathcal{A}^2bu^2 \pm \sqrt{4\alpha^2\mathcal{A}^2u^2(a^2 - b^2)(\mathcal{B} - 5\mathcal{A})^2 + (\mathcal{B} - 5\mathcal{A})^4}}{4a^2\alpha^2\mathcal{A}^2u^2 + (\mathcal{B} - 5\mathcal{A})^2}.$$

### 3.– Dirac eqn. with generalized gravitational interaction

#### First excited state and the associated wave function

For  $n = 1$ , the first excited-state energy,  $\epsilon_1$ , is given by

$$\epsilon_1 = \frac{-4a\alpha^2 \mathcal{A}^2 b u^2 \pm \sqrt{4\alpha^2 \mathcal{A}^2 u^2 (a^2 - b^2)(\mathcal{B} - 5\mathcal{A})^2 + (\mathcal{B} - 5\mathcal{A})^4}}{4a^2\alpha^2 \mathcal{A}^2 u^2 + (\mathcal{B} - 5\mathcal{A})^2}.$$

The associated wave function is  $\rho_{1,1}(r) \propto (r - r_1) e^{\Delta(r)}$ , with

$$\Delta(r) = \left( \frac{3}{2} + \frac{\alpha w(b - aC)(2\alpha u(b - aC) - 3S)}{2\sigma S^2} \right) \ln r - \frac{\sigma}{r^2} \left( \frac{rv}{w} + \frac{1}{2} \right) - \sqrt{\frac{1 - \epsilon_1^2}{\alpha^2}} r,$$

The potential parameters  $v$  and  $w$  are given in terms of the Coulombic coefficient  $u$  and  $a, b, C, S, \alpha, \epsilon_0$ .



### 3.– Dirac eqn. with generalized gravitational interaction

#### First excited state and the associated wave function

For  $n = 1$ , the first excited-state energy,  $\epsilon_1$ , is given by

$$\epsilon_1 = \frac{-4a\alpha^2 \mathcal{A}^2 bu^2 \pm \sqrt{4\alpha^2 \mathcal{A}^2 u^2 (a^2 - b^2)(\mathcal{B} - 5\mathcal{A})^2 + (\mathcal{B} - 5\mathcal{A})^4}}{4a^2\alpha^2 \mathcal{A}^2 u^2 + (\mathcal{B} - 5\mathcal{A})^2}.$$

The associated wave function is  $\rho_{1,1}(r) \propto (r - r_1) e^{\Delta(r)}$ , with

$$\Delta(r) = \left( \frac{3}{2} + \frac{\alpha w(b - aC)(2\alpha u(b - aC) - 3S)}{2\sigma S^2} \right) \ln r - \frac{\sigma}{r^2} \left( \frac{rv}{w} + \frac{1}{2} \right) - \sqrt{\frac{1 - \epsilon_1^2}{\alpha^2}} r,$$

The potential parameters  $v$  and  $w$  are given in terms of the Coulombic coefficient  $u$  and  $a, b, C, S, \alpha, \epsilon_0$ .

The special case of the Coulomb interaction  $v = w = 0$  can be obtained and compared the existing literature.

# Final Announcement:

# Final Announcement:



A conference in honor of Prof. J. NEGRO

# Final Announcement:



## A conference in honor of Prof. J. NEGRO

**XIth Int'l Workshop NEW CHALLENGES IN QUANTUM MECHANICS:  
Graphene, supersymmetry and mathematical physics**

**Valladolid (Spain), January 15-17, 2025**

# Final Announcement:



## A conference in honor of Prof. J. NEGRO

**XIth Int'l Workshop NEW CHALLENGES IN QUANTUM MECHANICS:  
Graphene, supersymmetry and mathematical physics**

**Valladolid (Spain), January 15-17, 2025**

### SCIENTIFIC TOPICS:

Special focus on the topics in which J. Negro has been working for years: coherent states, integrability and superintegrability, algebraic methods, continuous and discrete symmetries, quantum optics, quantum many-body systems, exact solutions in physics,...

# Final Announcement:



## A conference in honor of Prof. J. NEGRO

**XIth Int'l Workshop NEW CHALLENGES IN QUANTUM MECHANICS:  
Graphene, supersymmetry and mathematical physics**

**Valladolid (Spain), January 15-17, 2025**

### SCIENTIFIC TOPICS:

Special focus on the topics in which J. Negro has been working for years: coherent states, integrability and superintegrability, algebraic methods, continuous and discrete symmetries, quantum optics, quantum many-body systems, exact solutions in physics,...

**You are welcome to participate!**

**See you next January in Valladolid!**