

# Explicit $n$ -particle harmonic oscillator states

Hubert de Guise  
Lakehead University

and

Noah Kakekaspan  
now at McGill University



# Context

- In constructing multiparticle states, the permutation group is often essential.

# Context

- In constructing multiparticle states, the permutation group is often essential.
- The combination of two spin- $\frac{1}{2}$  systems gives:

# Context

- In constructing multiparticle states, the permutation group is often essential.
- The combination of two spin- $\frac{1}{2}$  systems gives:  
one *antisymmetric* singlet  $S = 0$  and

# Context

- In constructing multiparticle states, the permutation group is often essential.
- The combination of two spin- $\frac{1}{2}$  systems gives:  
one *antisymmetric* singlet  $S = 0$  and  
one *symmetric* triplet  $S = 1$ .

# Context

- In constructing multiparticle states, the permutation group is often essential.
- The combination of two spin- $\frac{1}{2}$  systems gives:  
one *antisymmetric* singlet  $S = 0$  and  
one *symmetric* triplet  $S = 1$ .
- The permutation symmetry uniquely identifies  $S$  when taking multiple copies of spin- $1/2$  states.

# Context

- In constructing multiparticle states, the permutation group is often essential.
- The combination of two spin- $\frac{1}{2}$  systems gives:  
one *antisymmetric* singlet  $S = 0$  and  
one *symmetric* triplet  $S = 1$ .
- The permutation symmetry uniquely identifies  $S$  when taking multiple copies of spin-1/2 states.
- In combining single-particle harmonic oscillator states of the same parity ( $\mathfrak{su}(1, 1) \sim \mathfrak{sp}(2, \mathbb{R})$  states), the permutation group is **not enough**.

# Promise

- Construct many-particle  $\mathfrak{su}(1,1)$  states “easily”,



# Promise

- Construct many-particle  $\mathfrak{su}(1, 1)$  states “easily”,
- Construct *some* many-particle  $\mathfrak{sp}(4, \mathbb{R})$  states “easily”

# Promise

- Construct many-particle  $\mathfrak{su}(1, 1)$  states “easily”,
- Construct *some* many-particle  $\mathfrak{sp}(4, \mathbb{R})$  states “easily”
- have an idea of where the permutation symmetry is hiding.

## $\mathfrak{su}(1,1)$ oscillator states

- The  $\mathfrak{su}(1,1) \sim \mathfrak{sp}(2, \mathbb{R})$  operators satisfy

$$[K_0, K_{\pm}] = \pm K_{\pm}, \quad [K_+, K_-] = -K_0.$$

# $\mathfrak{su}(1,1)$ oscillator states

- The  $\mathfrak{su}(1,1) \sim \mathfrak{sp}(2, \mathbb{R})$  operators satisfy

$$[K_0, K_{\pm}] = \pm K_{\pm}, \quad [K_+, K_-] = -K_0.$$

- (One particle) oscillator representation:

$$K_+ = \frac{1}{2} \hat{a}^\dagger \hat{a}^\dagger, \quad K_- = \frac{1}{2} \hat{a} \hat{a}, \quad K_0 = \frac{1}{2} (\hat{a}^\dagger \hat{a} + \frac{1}{2})$$

$$\begin{array}{c} \vdots \\ \text{---} |6\rangle \\ \text{---} |4\rangle \\ \text{---} |2\rangle \\ \text{---} |0\rangle \end{array}$$

# $\mathfrak{su}(1,1)$ oscillator states

- The  $\mathfrak{su}(1,1) \sim \mathfrak{sp}(2, \mathbb{R})$  operators satisfy

$$[K_0, K_{\pm}] = \pm K_{\pm}, \quad [K_+, K_-] = -K_0.$$

- (One particle) oscillator representation:

$$K_+ = \frac{1}{2} \hat{a}^\dagger \hat{a}^\dagger, \quad K_- = \frac{1}{2} \hat{a} \hat{a}, \quad K_0 = \frac{1}{2} (\hat{a}^\dagger \hat{a} + \frac{1}{2})$$

- The bottom state  $|0\rangle$  satisfies:

$$K_- |0\rangle = 0$$

$$\begin{array}{c} \vdots \\ \text{---} |6\rangle \\ \text{---} |4\rangle \\ \text{---} |2\rangle \\ \text{---} |0\rangle \end{array}$$

# $\mathfrak{su}(1,1)$ oscillator states

- The  $\mathfrak{su}(1,1) \sim \mathfrak{sp}(2, \mathbb{R})$  operators satisfy

$$[K_0, K_{\pm}] = \pm K_{\pm}, \quad [K_+, K_-] = -K_0.$$

- (One particle) oscillator representation:

$$K_+ = \frac{1}{2} \hat{a}^\dagger \hat{a}^\dagger, \quad K_- = \frac{1}{2} \hat{a} \hat{a}, \quad K_0 = \frac{1}{2} (\hat{a}^\dagger \hat{a} + \frac{1}{2})$$

- The bottom state  $|0\rangle$  satisfies:

$$K_- |0\rangle = 0$$

- Natural for boson systems

IOP Publishing

Journal of Physics A: Mathematical and Theoretical

J. Phys. A: Math. Theor. **47** (2014) 275202 (9pp)

[doi:10.1088/1751-8113/47/27/275202](https://doi.org/10.1088/1751-8113/47/27/275202)

## Where do bosons actually belong?

A Marzuoli<sup>1,2</sup>, F A Raffa<sup>3</sup> and M Rasetti<sup>3,4</sup>

## Combining two su(1,1) systems

- $K_- = \frac{1}{2} (\hat{a}_1 \hat{a}_1 + \hat{a}_2 \hat{a}_2)$  ,  $K_+ = K_-^\dagger$  ,  $K_0 = \frac{1}{2} (\hat{a}_1^\dagger \hat{a}_1 + \hat{a}_2^\dagger \hat{a}_2 + 1)$

## Combining two $\text{su}(1,1)$ systems

- $K_- = \frac{1}{2} (\hat{a}_1 \hat{a}_1 + \hat{a}_2 \hat{a}_2)$  ,  $K_+ = K_-^\dagger$  ,  $K_0 = \frac{1}{2} (\hat{a}_1^\dagger \hat{a}_1 + \hat{a}_2^\dagger \hat{a}_2 + 1)$
- All states in a tower have the same permutation symmetry.



# Combining two su(1,1) systems

- $K_- = \frac{1}{2} (\hat{a}_1 \hat{a}_1 + \hat{a}_2 \hat{a}_2)$  ,  $K_+ = K_-^\dagger$  ,  $K_0 = \frac{1}{2} (\hat{a}_1^\dagger \hat{a}_1 + \hat{a}_2^\dagger \hat{a}_2 + 1)$
- All states in a tower have the same permutation symmetry.
- There are infinitely many towers of symmetric states.

⋮

—

—  $|4\rangle|0\rangle + \sqrt{\frac{2}{3}}|2\rangle|2\rangle + |0\rangle|4\rangle$

—  $|2\rangle|0\rangle + |0\rangle|2\rangle$

—  $|0\rangle|0\rangle$

⋮

—

—  $|4\rangle|0\rangle - \sqrt{6}|2\rangle|2\rangle + |0\rangle|4\rangle$

# Combining two su(1,1) systems

- $K_- = \frac{1}{2} (\hat{a}_1 \hat{a}_1 + \hat{a}_2 \hat{a}_2)$ ,  $K_+ = K_-^\dagger$ ,  $K_0 = \frac{1}{2} (\hat{a}_1^\dagger \hat{a}_1 + \hat{a}_2^\dagger \hat{a}_2 + 1)$
- All states in a tower have the same permutation symmetry.
- There are infinitely many towers of symmetric states.

⋮

—

—  $|4\rangle|0\rangle - |0\rangle|4\rangle$

—  $|2\rangle|0\rangle - |0\rangle|2\rangle$

- There are infinitely many towers of antisymmetric states.

# Combining two su(1,1) systems

- $K_- = \frac{1}{2} (\hat{a}_1 \hat{a}_1 + \hat{a}_2 \hat{a}_2)$ ,  $K_+ = K_-^\dagger$ ,  $K_0 = \frac{1}{2} (\hat{a}_1^\dagger \hat{a}_1 + \hat{a}_2^\dagger \hat{a}_2 + 1)$
- All states in a tower have the same permutation symmetry.
- There are infinitely many towers of symmetric states.

$\vdots$	$\vdots$	$\vdots$
—	—	—
— $ 4\rangle 0\rangle + \sqrt{\frac{2}{3}} 2\rangle 2\rangle +  0\rangle 4\rangle$	— $ 4\rangle 0\rangle -  0\rangle 4\rangle$	— $ 4\rangle 0\rangle - \sqrt{6} 2\rangle 2\rangle +  0\rangle 4\rangle$
— $ 2\rangle 0\rangle +  0\rangle 2\rangle$	— $ 2\rangle 0\rangle -  0\rangle 2\rangle$	
— $ 0\rangle 0\rangle$		

- Clearly permutation symmetry is not enough to identify the tower of states.

# The Laplacian approach

$\vdots$	$\vdots$	$\vdots$
—	—	—
— $ 4\rangle 0\rangle + \sqrt{\frac{2}{3}} 2\rangle 2\rangle +  0\rangle 4\rangle$	— $ 4\rangle 0\rangle -  0\rangle 4\rangle$	— $ 4\rangle 0\rangle - \sqrt{6} 2\rangle 2\rangle +  0\rangle 4\rangle$
— $ 2\rangle 0\rangle +  0\rangle 2\rangle$	— $ 2\rangle 0\rangle -  0\rangle 2\rangle$	
— $ 0\rangle 0\rangle$		

# The Laplacian approach

$\vdots$	$\vdots$	$\vdots$
—	—	—
— $ 4\rangle 0\rangle + \sqrt{\frac{2}{3}} 2\rangle 2\rangle +  0\rangle 4\rangle$	— $ 4\rangle 0\rangle -  0\rangle 4\rangle$	— $ 4\rangle 0\rangle - \sqrt{6} 2\rangle 2\rangle +  0\rangle 4\rangle$
— $ 2\rangle 0\rangle +  0\rangle 2\rangle$	— $ 2\rangle 0\rangle -  0\rangle 2\rangle$	
— $ 0\rangle 0\rangle$		

- The 3 bottom states satisfy

# The Laplacian approach

$\vdots$	$\vdots$	$\vdots$
—	—	—
— $ 4\rangle 0\rangle + \sqrt{\frac{2}{3}} 2\rangle 2\rangle +  0\rangle 4\rangle$	— $ 4\rangle 0\rangle -  0\rangle 4\rangle$	— $ 4\rangle 0\rangle - \sqrt{6} 2\rangle 2\rangle +  0\rangle 4\rangle$
— $ 2\rangle 0\rangle +  0\rangle 2\rangle$	— $ 2\rangle 0\rangle -  0\rangle 2\rangle$	
— $ 0\rangle 0\rangle$		

- The 3 bottom states satisfy

$$\begin{aligned}
 K_-|0\rangle|0\rangle &= 0, \\
 K_- (|2\rangle|0\rangle - |0\rangle|2\rangle) &= 0, \\
 K_- (|4\rangle|0\rangle - \sqrt{6}|2\rangle|2\rangle + |0\rangle|4\rangle) &= 0
 \end{aligned}$$

# The Laplacian approach

- Write

$$\hat{a}_i \mapsto \frac{\partial}{\partial x_i}, \quad \hat{a}_j^\dagger \mapsto x_j \quad \Rightarrow \quad K_- \mapsto \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2}$$

# The Laplacian approach

- Write

$$\hat{a}_i \mapsto \frac{\partial}{\partial x_i}, \quad \hat{a}_j^\dagger \mapsto x_j \quad \Rightarrow \quad K_- \mapsto \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2}$$

- The bottom states map to polynomials:

$$|0\rangle|0\rangle \mapsto f_0 = 1,$$

$$|2\rangle|0\rangle - |0\rangle|2\rangle \mapsto f_2 = x_1^2 - x_2^2,$$

$$|4\rangle|0\rangle - \sqrt{6}|2\rangle|2\rangle + |0\rangle|4\rangle \mapsto f_4 = x_1^4 - 6x_1^2x_2^2 + x_2^4$$



# The Laplacian approach

- Write

$$\hat{a}_i \mapsto \frac{\partial}{\partial x_i}, \quad \hat{a}_j^\dagger \mapsto x_j \quad \Rightarrow \quad K_- \mapsto \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2}$$

- The bottom states map to polynomials:

$$|0\rangle|0\rangle \mapsto f_0 = 1,$$

$$|2\rangle|0\rangle - |0\rangle|2\rangle \mapsto f_2 = x_1^2 - x_2^2,$$

$$|4\rangle|0\rangle - \sqrt{6}|2\rangle|2\rangle + |0\rangle|4\rangle \mapsto f_4 = x_1^4 - 6x_1^2x_2^2 + x_2^4$$

- Now we have

$$K_- f_k \mapsto \nabla^2 f_k = 0.$$

# The Laplacian approach

- Write

$$\hat{a}_i \mapsto \frac{\partial}{\partial x_i}, \quad \hat{a}_j^\dagger \mapsto x_j \quad \Rightarrow \quad K_- \mapsto \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2}$$

- The bottom states map to polynomials:

$$|0\rangle|0\rangle \mapsto f_0 = 1,$$

$$|2\rangle|0\rangle - |0\rangle|2\rangle \mapsto f_2 = x_1^2 - x_2^2,$$

$$|4\rangle|0\rangle - \sqrt{6}|2\rangle|2\rangle + |0\rangle|4\rangle \mapsto f_4 = x_1^4 - 6x_1^2x_2^2 + x_2^4$$

- Now we have

$$K_- f_k \mapsto \nabla^2 f_k = 0.$$

- The  $f_k$  are two-dimensional harmonic functions.

# The Laplacian approach

- Solutions  $f_k$  have a generating function:

# The Laplacian approach

- Solutions  $f_k$  have a generating function:

$$H_2^+(x_1, x_2) = e^{(x_1 + ix_2)h_2}$$

# The Laplacian approach

- Solutions  $f_k$  have a generating function:

$$H_2^+(x_1, x_2) = e^{(x_1 + ix_2)h_2} \approx 1 + h_2(x_1 + ix_2) + h_2^2 \frac{1}{2}(x_1 + ix_2)^2 + \dots$$

# The Laplacian approach

- Solutions  $f_k$  have a generating function:

$$H_2^+(x_1, x_2) = e^{(x_1 + ix_2)h_2} \approx 1 + h_2(x_1 + ix_2) + h_2^2 \frac{1}{2}(x_1 + ix_2)^2 + \dots$$

- In the expansion of  $H_2^+(x_1, x_2)$ , distinct powers of  $h_2$  give distinct functions  $f_k$  that satisfy  $\nabla^2 f_k = 0$ .

# The Laplacian approach

- Solutions  $f_k$  have a generating function:

$$H_2^+(x_1, x_2) = e^{(x_1 + ix_2)h_2} \approx 1 + h_2(x_1 + ix_2) + h_2^2 \frac{1}{2}(x_1 + ix_2)^2 + \dots$$

- In the expansion of  $H_2^+(x_1, x_2)$ , distinct powers of  $h_2$  give distinct functions  $f_k$  that satisfy  $\nabla^2 f_k = 0$ .
- Only even numbers of excitations are possible so even powers of  $x_1$  or  $x_2$  must be kept:

$$1 + \frac{1}{2}h_2^2(x_1^2 - x_2^2) + \frac{1}{24}h_2^4(x_1^4 - 6x_1^2x_2^2 + x_2^4) + \dots$$

# The Laplacian approach

- For three particles:

$$\nabla^2 = \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \frac{\partial^2}{\partial x_3^2} = 0.$$



# The Laplacian approach

- For three particles:

$$\nabla^2 = \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \frac{\partial^2}{\partial x_3^2} = 0.$$

- There's also a generating function for the solutions to this.

Adv. Appl. Clifford Algebras 24 (2014), 995–1004  
© 2014 Springer Basel  
0188-7009/040995-10  
*published online* August 31, 2014  
DOI 10.1007/s00006-014-0495-8

Advances in  
Applied Clifford Algebras

## Generating Functions for Spherical Harmonics and Spherical Monogenics

P. Cerejeiras, U. Kähler and R. Lávička\*

# Connection with spherical harmonics

- Again keeping only terms in even powers of  $x_i$ :

$$1 + \frac{1}{2}(x_1^2 - x_2^2) + \frac{1}{2}(x_1^2 + x_2^2 - 2x_3^2)$$

# Connection with spherical harmonics

- Again keeping only terms in even powers of  $x_i$ :

$$1 + \frac{1}{2}(x_1^2 - x_2^2) + \frac{1}{2}(x_1^2 + x_2^2 - 2x_3^2)$$

- Thus:

$$|0\rangle|0\rangle|0\rangle, \quad |2\rangle|0\rangle|0\rangle - |0\rangle|2\rangle|0\rangle, \quad |2\rangle|0\rangle|0\rangle + |0\rangle|2\rangle|0\rangle - 2|0\rangle|0\rangle|2\rangle$$

are all at the bottom of separate 3-particle towers.

# Connection with spherical harmonics

- Again keeping only terms in even powers of  $x_i$ :

$$1 + \frac{1}{2}(x_1^2 - x_2^2) + \frac{1}{2}(x_1^2 + x_2^2 - 2x_3^2)$$

- Thus:

$$|0\rangle|0\rangle|0\rangle, \quad |2\rangle|0\rangle|0\rangle - |0\rangle|2\rangle|0\rangle, \quad |2\rangle|0\rangle|0\rangle + |0\rangle|2\rangle|0\rangle - 2|0\rangle|0\rangle|2\rangle$$

are all at the bottom of separate 3-particle towers.

- of course

# Connection with spherical harmonics

- Again keeping only terms in even powers of  $x_i$ :

$$1 + \frac{1}{2}(x_1^2 - x_2^2) + \frac{1}{2}(x_1^2 + x_2^2 - 2x_3^2)$$

- Thus:

$$|0\rangle|0\rangle|0\rangle, \quad |2\rangle|0\rangle|0\rangle - |0\rangle|2\rangle|0\rangle, \quad |2\rangle|0\rangle|0\rangle + |0\rangle|2\rangle|0\rangle - 2|0\rangle|0\rangle|2\rangle$$

are all at the bottom of separate 3-particle towers.

- of course

$$r^2(Y_2^{-2}(\theta, \varphi) + Y_2^2(\theta, \varphi)) \sim x_1^2 - x_2^2$$

# Connection with spherical harmonics

- Again keeping only terms in even powers of  $x_i$ :

$$1 + \frac{1}{2}(x_1^2 - x_2^2) + \frac{1}{2}(x_1^2 + x_2^2 - 2x_3^2)$$

- Thus:

$$|0\rangle|0\rangle|0\rangle, \quad |2\rangle|0\rangle|0\rangle - |0\rangle|2\rangle|0\rangle, \quad |2\rangle|0\rangle|0\rangle + |0\rangle|2\rangle|0\rangle - 2|0\rangle|0\rangle|2\rangle$$

are all at the bottom of separate 3-particle towers.

- of course

$$\begin{aligned} r^2(Y_2^{-2}(\theta, \varphi) + Y_2^2(\theta, \varphi)) &\sim x_1^2 - x_2^2 \\ r^2 Y_2^0(\theta, \varphi) &\sim x_1^2 + x_2^2 - 2x_3^2 \end{aligned}$$

# Connection with spherical harmonics

- Again keeping only terms in even powers of  $x_i$ :

$$1 + \frac{1}{2}(x_1^2 - x_2^2) + \frac{1}{2}(x_1^2 + x_2^2 - 2x_3^2)$$

- Thus:

$$|0\rangle|0\rangle|0\rangle, \quad |2\rangle|0\rangle|0\rangle - |0\rangle|2\rangle|0\rangle, \quad |2\rangle|0\rangle|0\rangle + |0\rangle|2\rangle|0\rangle - 2|0\rangle|0\rangle|2\rangle$$

are all at the bottom of separate 3-particle towers.

- of course

$$\begin{aligned} r^2(Y_2^{-2}(\theta, \varphi) + Y_2^2(\theta, \varphi)) &\sim x_1^2 - x_2^2 \\ r^2 Y_2^0(\theta, \varphi) &\sim x_1^2 + x_2^2 - 2x_3^2 \end{aligned}$$

- There is in fact a symmetry to the solutions: angular momentum plus parity.

## Three particles and $O(3)$

- For 3 particles we have:

$$K_+ = \frac{1}{2} \left( \hat{a}_1^\dagger \hat{a}_1^\dagger + \hat{a}_2^\dagger \hat{a}_2^\dagger + \hat{a}_3^\dagger \hat{a}_3^\dagger \right), \quad K_- = K_+^\dagger,$$
$$K_0 = \frac{1}{2} \left( \hat{a}_1^\dagger \hat{a}_1 + \hat{a}_2^\dagger \hat{a}_2 + \hat{a}_3^\dagger \hat{a}_3 + \frac{3}{2} \right)$$



# Three particles and $O(3)$

- For 3 particles we have:

$$K_+ = \frac{1}{2} \left( \hat{a}_1^\dagger \hat{a}_1^\dagger + \hat{a}_2^\dagger \hat{a}_2^\dagger + \hat{a}_3^\dagger \hat{a}_3^\dagger \right), \quad K_- = K_+^\dagger,$$
$$K_0 = \frac{1}{2} \left( \hat{a}_1^\dagger \hat{a}_1 + \hat{a}_2^\dagger \hat{a}_2 + \hat{a}_3^\dagger \hat{a}_3 + \frac{3}{2} \right)$$

- Introduce

$$\hat{L}_{12} = i \left( \hat{a}_1^\dagger \hat{a}_2 - \hat{a}_2^\dagger \hat{a}_1 \right) \rightarrow \hat{L}_z, \quad \hat{L}_{23} = i \left( \hat{a}_2^\dagger \hat{a}_3 - \hat{a}_3^\dagger \hat{a}_2 \right) \rightarrow \hat{L}_x,$$
$$\hat{L}_{31} = i \left( \hat{a}_3^\dagger \hat{a}_1 - \hat{a}_1^\dagger \hat{a}_3 \right) \rightarrow \hat{L}_y.$$

# Three particles and $O(3)$

- For 3 particles we have:

$$K_+ = \frac{1}{2} \left( \hat{a}_1^\dagger \hat{a}_1^\dagger + \hat{a}_2^\dagger \hat{a}_2^\dagger + \hat{a}_3^\dagger \hat{a}_3^\dagger \right), \quad K_- = K_+^\dagger,$$
$$K_0 = \frac{1}{2} \left( \hat{a}_1^\dagger \hat{a}_1 + \hat{a}_2^\dagger \hat{a}_2 + \hat{a}_3^\dagger \hat{a}_3 + \frac{3}{2} \right)$$

- Introduce

$$\hat{L}_{12} = i \left( \hat{a}_1^\dagger \hat{a}_2 - \hat{a}_2^\dagger \hat{a}_1 \right) \rightarrow \hat{L}_z, \quad \hat{L}_{23} = i \left( \hat{a}_2^\dagger \hat{a}_3 - \hat{a}_3^\dagger \hat{a}_2 \right) \rightarrow \hat{L}_x,$$
$$\hat{L}_{31} = i \left( \hat{a}_3^\dagger \hat{a}_1 - \hat{a}_1^\dagger \hat{a}_3 \right) \rightarrow \hat{L}_y.$$

- Of course  $[\hat{L}_j, \hat{L}_k] = i\epsilon_{jkl} \hat{L}_l$

# Three particles and $O(3)$

- For 3 particles we have:

$$K_+ = \frac{1}{2} \left( \hat{a}_1^\dagger \hat{a}_1^\dagger + \hat{a}_2^\dagger \hat{a}_2^\dagger + \hat{a}_3^\dagger \hat{a}_3^\dagger \right), \quad K_- = K_+^\dagger,$$
$$K_0 = \frac{1}{2} \left( \hat{a}_1^\dagger \hat{a}_1 + \hat{a}_2^\dagger \hat{a}_2 + \hat{a}_3^\dagger \hat{a}_3 + \frac{3}{2} \right)$$

- Introduce

$$\hat{L}_{12} = i \left( \hat{a}_1^\dagger \hat{a}_2 - \hat{a}_2^\dagger \hat{a}_1 \right) \rightarrow \hat{L}_z, \quad \hat{L}_{23} = i \left( \hat{a}_2^\dagger \hat{a}_3 - \hat{a}_3^\dagger \hat{a}_2 \right) \rightarrow \hat{L}_x,$$
$$\hat{L}_{31} = i \left( \hat{a}_3^\dagger \hat{a}_1 - \hat{a}_1^\dagger \hat{a}_3 \right) \rightarrow \hat{L}_y.$$

- Of course  $[\hat{L}_j, \hat{L}_k] = i\epsilon_{jkl} \hat{L}_l$
- Also  $[\hat{L}_j, K_q] = 0$ .

## Three particles and $O(3)$

- Thus, all states in a tower are also "good angular momentum states", *i.e.* they have good quantum numbers  $\ell$  and  $|m|$ .

## Three particles and $O(3)$

- Thus, all states in a tower are also "good angular momentum states", *i.e.* they have good quantum numbers  $\ell$  and  $|m|$ .
- Because of parity, states are combo of  $|m|$  states with  $m$  even.

## Three particles and $O(3)$

- Thus, all states in a tower are also "good angular momentum states", *i.e.* they have good quantum numbers  $\ell$  and  $|m|$ .
- Because of parity, states are combo of  $|m|$  states with  $m$  even.
- Recall

$$K_+ = a_1^\dagger a_1^\dagger + a_2^\dagger a_2^\dagger + a_3^\dagger a_3^\dagger$$

## Three particles and $O(3)$

- Thus, all states in a tower are also "good angular momentum states", *i.e.* they have good quantum numbers  $\ell$  and  $|m|$ .
- Because of parity, states are combo of  $|m|$  states with  $m$  even.
- Recall

$$K_+ = a_1^\dagger a_1^\dagger + a_2^\dagger a_2^\dagger + a_3^\dagger a_3^\dagger \quad \mapsto x_1^2 + x_2^2 + x_3^2 = r^2$$

# Three particles and $O(3)$

- Thus, all states in a tower are also "good angular momentum states", *i.e.* they have good quantum numbers  $\ell$  and  $|m|$ .
- Because of parity, states are combo of  $|m|$  states with  $m$  even.
- Recall

$$K_+ = a_1^\dagger a_1^\dagger + a_2^\dagger a_2^\dagger + a_3^\dagger a_3^\dagger \quad \mapsto x_1^2 + x_2^2 + x_3^2 = r^2$$

- The states  $|\ell, m; n\rangle$  in  $\ell = 0, m = 0$  tower:

$$Y_0^0(\theta, \varphi) = 1 \quad \sim |0\rangle|0\rangle|0\rangle \quad = |00; 0\rangle$$

$$r^2 Y_0^0(\theta, \varphi) = x_1^2 + x_2^2 + x_3^2 \quad \sim |2\rangle|0\rangle|0\rangle + |0\rangle|2\rangle|0\rangle + |0\rangle|0\rangle|2\rangle = |00; 2\rangle$$

$$r^4 Y_0^0(\theta, \varphi) = (x_1^2 + x_2^2 + x_3^2)^2 \quad = |00; 4\rangle$$



# Three particles and $O(3)$

- The  $\ell = 4, m = 0$  tower are built on:

$$r^4 Y_4^0(\theta, \varphi) \sim |40; 4\rangle = |4\rangle|0\rangle|0\rangle + \sqrt{\frac{2}{3}}|2\rangle|2\rangle|0\rangle - 4\sqrt{\frac{2}{3}}|2\rangle|0\rangle|2\rangle + \frac{8}{3}|0\rangle|0\rangle|4\rangle$$

# Three particles and $O(3)$

- The  $\ell = 4, m = 0$  tower are built on:

$$r^4 Y_4^0(\theta, \varphi) \sim |40; 4\rangle = |4\rangle|0\rangle|0\rangle + \sqrt{\frac{2}{3}}|2\rangle|2\rangle|0\rangle - 4\sqrt{\frac{2}{3}}|2\rangle|0\rangle|2\rangle + \frac{8}{3}|0\rangle|0\rangle|4\rangle$$

- We also have  $\ell = 4, |m| = 2$  and  $|m| = 4$  towers built on:

$$\begin{aligned} r^4(Y_4^2(\theta, \varphi) + Y_4^{-2}(\theta, \varphi)) \sim |42; 4\rangle \sim & |4\rangle|0\rangle|0\rangle - \sqrt{6}|2\rangle|0\rangle|2\rangle \\ & - |0\rangle|4\rangle|0\rangle + \sqrt{6}|0\rangle|2\rangle|2\rangle \end{aligned}$$

$$r^4(Y_4^4(\theta, \varphi) + Y_4^{-4}(\theta, \varphi)) \sim |44; 4\rangle \sim |4\rangle|0\rangle|0\rangle - \sqrt{6}|2\rangle|2\rangle|0\rangle + |0\rangle|4\rangle|0\rangle$$

# Where did the permutation symmetry go?

# Where did the permutation symmetry go?

- The combination  $|40; 4\rangle + \frac{5}{3}|44; 4\rangle$  is proportional to

$$|0\rangle|0\rangle|4\rangle + |0\rangle|4\rangle|0\rangle + |4\rangle|0\rangle|0\rangle - \sqrt{\frac{3}{2}}(|2\rangle|2\rangle|0\rangle + |2\rangle|0\rangle|2\rangle + |0\rangle|2\rangle|2\rangle)$$

and is fully symmetric, like the  $|00; 0\rangle$  state.

# Where did the permutation symmetry go?

- The combination  $|40; 4\rangle + \frac{5}{3}|44; 4\rangle$  is proportional to

$$|0\rangle|0\rangle|4\rangle + |0\rangle|4\rangle|0\rangle + |4\rangle|0\rangle|0\rangle - \sqrt{\frac{3}{2}}(|2\rangle|2\rangle|0\rangle + |2\rangle|0\rangle|2\rangle + |0\rangle|2\rangle|2\rangle)$$

and is fully symmetric, like the  $|00; 0\rangle$  state.

- There are two remaining states with are partially symmetric.

# Where did the permutation symmetry go?

- The combination  $|40; 4\rangle + \frac{5}{3}|44; 4\rangle$  is proportional to

$$|0\rangle|0\rangle|4\rangle + |0\rangle|4\rangle|0\rangle + |4\rangle|0\rangle|0\rangle - \sqrt{\frac{3}{2}}(|2\rangle|2\rangle|0\rangle + |2\rangle|0\rangle|2\rangle + |0\rangle|2\rangle|2\rangle)$$

and is fully symmetric, like the  $|00; 0\rangle$  state.

- There are two remaining states with are partially symmetric.  
One goes to a linear combination of the other two under permutation.

# Where did the permutation symmetry go?

- The combination  $|40; 4\rangle + \frac{5}{3}|44; 4\rangle$  is proportional to

$$|0\rangle|0\rangle|4\rangle + |0\rangle|4\rangle|0\rangle + |4\rangle|0\rangle|0\rangle - \sqrt{\frac{3}{2}}(|2\rangle|2\rangle|0\rangle + |2\rangle|0\rangle|2\rangle + |0\rangle|2\rangle|2\rangle)$$

and is fully symmetric, like the  $|00; 0\rangle$  state.

- There are two remaining states with are partially symmetric.  
One goes to a linear combination of the other two under permutation.
- The permutation group appears as a subgroup of  $O(3)$ :

# Where did the permutation symmetry go?

- The combination  $|40; 4\rangle + \frac{5}{3}|44; 4\rangle$  is proportional to

$$|0\rangle|0\rangle|4\rangle + |0\rangle|4\rangle|0\rangle + |4\rangle|0\rangle|0\rangle - \sqrt{\frac{3}{2}}(|2\rangle|2\rangle|0\rangle + |2\rangle|0\rangle|2\rangle + |0\rangle|2\rangle|2\rangle)$$

and is fully symmetric, like the  $|00; 0\rangle$  state.

- There are two remaining states with are partially symmetric. One goes to a linear combination of the other two under permutation.
- The permutation group appears as a subgroup of  $O(3)$ :

$$P_{12} = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \text{ is an orthogonal matrix: } P_{12}^T = P_{12}^{-1}$$



## More than 3 particles

- The strategy works for  $d \geq 3$  particles but...

# More than 3 particles

- The strategy works for  $d \geq 3$  particles but...
  - The generating function becomes complicated.

# More than 3 particles

- The strategy works for  $d \geq 3$  particles but...
  - The generating function becomes complicated.
  - Other methods use hyperspherical coordinates.

# More than 3 particles

- The strategy works for  $d \geq 3$  particles but...
  - The generating function becomes complicated.
  - Other methods use hyperspherical coordinates.
  - Tensor operator methods

# Single-particle irrep of $sp(4, \mathbb{R})$

- We have the generators

$$A_{ij} = a_i^\dagger a_j^\dagger, \quad i = 1, 2,$$

$$C_{ij} = \frac{1}{2} \left( a_i^\dagger a_j + a_j a_i^\dagger \right), \quad i = 1, 2,$$

$$B_{ij} = a_i a_j, \quad i = 1, 2.$$

# Single-particle irrep of $sp(4, \mathbb{R})$

- We have the generators

$$A_{ij} = a_i^\dagger a_j^\dagger, \quad i = 1, 2,$$

$$C_{ij} = \frac{1}{2} \left( a_i^\dagger a_j + a_j a_i^\dagger \right), \quad i = 1, 2,$$

$$B_{ij} = a_i a_j, \quad i = 1, 2.$$

- The set

$$C_{ij} = \frac{1}{2} \left( a_i^\dagger a_j + a_j a_i^\dagger \right), \quad i = 1, 2,$$

span a  $u(2)$  subalgebra.

# Single-particle irrep of $sp(4, \mathbb{R})$

- We have the generators

$$A_{ij} = a_i^\dagger a_j^\dagger, \quad i = 1, 2,$$

$$C_{ij} = \frac{1}{2} \left( a_i^\dagger a_j + a_j a_i^\dagger \right), \quad i = 1, 2,$$

$$B_{ij} = a_i a_j, \quad i = 1, 2.$$

- The set

$$C_{ij} = \frac{1}{2} \left( a_i^\dagger a_j + a_j a_i^\dagger \right), \quad i = 1, 2,$$

span a  $u(2)$  subalgebra.

- The operators  $\{A_{11}, A_{12}, A_{22}\}$  transform under the  $u(2)$  as states in the  $u(2)$  irrep  $(2, 0)$ .

# Single-particle irrep of $sp(4, \mathbb{R})$

- We have the generators

$$A_{ij} = a_i^\dagger a_j^\dagger, \quad i = 1, 2,$$

$$C_{ij} = \frac{1}{2} \left( a_i^\dagger a_j + a_j a_i^\dagger \right), \quad i = 1, 2,$$

$$B_{ij} = a_i a_j, \quad i = 1, 2.$$

- The set

$$C_{ij} = \frac{1}{2} \left( a_i^\dagger a_j + a_j a_i^\dagger \right), \quad i = 1, 2,$$

span a  $u(2)$  subalgebra.

- The operators  $\{A_{11}, A_{12}, A_{22}\}$  transform under the  $u(2)$  as states in the  $u(2)$  irrep  $(2, 0)$ .
- As per  $su(1, 1)$ , the idea is to start from the bottom of the tower and repeated act with powers of the  $A_{ij}$ :

$$A_{ij} A_{kl} \in [(2, 0) \otimes (2, 0)]_{\text{symm}}$$



# Single-particle irrep of $sp(4, \mathbb{R})$

- We have the generators

$$A_{ij} = a_i^\dagger a_j^\dagger, \quad i = 1, 2,$$

$$C_{ij} = \frac{1}{2} \left( a_i^\dagger a_j + a_j a_i^\dagger \right), \quad i = 1, 2,$$

$$B_{ij} = a_i a_j, \quad i = 1, 2.$$

- The set

$$C_{ij} = \frac{1}{2} \left( a_i^\dagger a_j + a_j a_i^\dagger \right), \quad i = 1, 2,$$

span a  $u(2)$  subalgebra.

- The operators  $\{A_{11}, A_{12}, A_{22}\}$  transform under the  $u(2)$  as states in the  $u(2)$  irrep  $(2, 0)$ .
- As per  $su(1, 1)$ , the idea is to start from the bottom of the tower and repeated act with powers of the  $A_{ij}$ :

$$A_{ij} A_{kl} \in [(2, 0) \otimes (2, 0)]_{\text{symm}}$$

- Because  $A_{ij}$ 's commute only the symmetric part of the tensor product is needed.

# Irreps of $sp(4, \mathbb{R})$

- The symmetric part of the repeated coupling of  $(2, 0)$  irreps contains states in the  $u(2)$  irreps

$$D = (00) + (20) + (40) + (22) + (60) + (42) + (22) + \dots$$

# Irreps of $sp(4, \mathbb{R})$

- The symmetric part of the repeated coupling of  $(2, 0)$  irreps contains states in the  $u(2)$  irreps

$$D = (00) + (20) + (40) + (22) + (60) + (42) + (22) + \dots$$

- the sum is over partitions of even integers having even-only parts.

J. Phys. A: Math. Gen. **18** (1985) 939-953. Printed in Great Britain

## **Unitary representations, branching rules and matrix elements for the non-compact symplectic groups**

D J Rowe<sup>†</sup>, B G Wybourne and P H Butler

Physics Department, University of Canterbury, Christchurch 1, New Zealand

# Irreps of $sp(4, \mathbb{R})$

- The symmetric part of the repeated coupling of  $(2, 0)$  irreps contains states in the  $u(2)$  irreps

$$D = (00) + (20) + (40) + (22) + (60) + (42) + (22) + \dots$$

- the sum is over partitions of even integers having even-only parts.
- These must in turn be coupled to the bottom state of each tower.

J. Phys. A: Math. Gen. **18** (1985) 939-953. Printed in Great Britain

## **Unitary representations, branching rules and matrix elements for the non-compact symplectic groups**

D J Rowe<sup>†</sup>, B G Wybourne and P H Butler

Physics Department, University of Canterbury, Christchurch 1, New Zealand

# Irreps of $sp(4, \mathbb{R})$

- The symmetric part of the repeated coupling of  $(2, 0)$  irreps contains states in the  $u(2)$  irreps

$$D = (00) + (20) + (40) + (22) + (60) + (42) + (22) + \dots$$

- the sum is over partitions of even integers having even-only parts.
- These must in turn be coupled to the bottom state of each tower.
- For single particles irrep of  $sp(4, \mathbb{R})$  this bottom state is carries the  $(0, 0)$  irrep so  $D$  lists the  $u(2)$  contents of the single particle irrep.

J. Phys. A: Math. Gen. **18** (1985) 939-953. Printed in Great Britain

## **Unitary representations, branching rules and matrix elements for the non-compact symplectic groups**

D J Rowe†, B G Wybourne and P H Butler

Physics Department, University of Canterbury, Christchurch 1, New Zealand

## Two-particle irreps of $sp(4, \mathbb{R})$

- We now have

$$A_{ij} = \sum_{k=1}^2 a_{ki}^{\dagger} a_{kj}^{\dagger},$$

$$C_{ij} = \sum_{k=1}^2 \frac{1}{2} \left( a_{ki}^{\dagger} a_{kj} + a_{kj} a_{ki}^{\dagger} \right),$$

$$B_{ij} = A_{ij}^{\dagger}.$$

# Two-particle irreps of $sp(4, \mathbb{R})$

- We now have

$$A_{ij} = \sum_{k=1}^2 a_{ki}^{\dagger} a_{kj}^{\dagger},$$

$$C_{ij} = \sum_{k=1}^2 \frac{1}{2} \left( a_{ki}^{\dagger} a_{kj} + a_{kj} a_{ki}^{\dagger} \right),$$

$$B_{ij} = A_{ij}^{\dagger}.$$

- The lowering operators

$$B_{11} \mapsto \frac{\partial^2}{\partial x_{11}^2} + \frac{\partial^2}{\partial x_{21}^2},$$

$$B_{12} \mapsto \frac{\partial^2}{\partial x_{11} \partial x_{12}} + \frac{\partial^2}{\partial x_{21} \partial x_{22}},$$

$$B_{22} \mapsto \frac{\partial^2}{\partial x_{12}^2} + \frac{\partial^2}{\partial x_{22}^2}.$$

## Two-particle irreps of $sp(4, \mathbb{R})$

- Not so easy to find the functions  $f$  *simultaneously* solutions to

$$\begin{aligned}\left(\frac{\partial^2}{\partial x_{11}^2} + \frac{\partial^2}{\partial x_{21}^2}\right) f &= 0, \\ \left(\frac{\partial^2}{\partial x_{11} \partial x_{12}} + \frac{\partial^2}{\partial x_{21} \partial x_{22}}\right) f &= 0 \\ \left(\frac{\partial^2}{\partial x_{12}^2} + \frac{\partial^2}{\partial x_{22}^2}\right) f &= 0.\end{aligned}$$



## Two-particle irreps of $sp(4, \mathbb{R})$

- Use  $O(2)$  by constructing the tensors

$$T^1 = \frac{1}{2} \left( a_{11}^\dagger - i a_{21}^\dagger \right), \quad V^1 = \frac{1}{2} \left( a_{12}^\dagger - i a_{22}^\dagger \right).$$

$a_{i\alpha}^\dagger$ : particle number  $i$ , mode number  $\alpha$ .

- Note that

**Clarify what is L12**

$$[L_{12}, T^1] = +1T^1, \quad [L_{12}, V^1] = +1V^1$$

## Two-particle irreps of $sp(4, \mathbb{R})$

- Use  $O(2)$  by constructing the tensors

$$T^1 = \frac{1}{2} \left( a_{11}^\dagger - i a_{21}^\dagger \right), \quad V^1 = \frac{1}{2} \left( a_{12}^\dagger - i a_{22}^\dagger \right).$$

$a_{i\alpha}^\dagger$ : particle number  $i$ , mode number  $\alpha$ .

- Note that

$$[L_{12}, T^1] = +1T^1, \quad [L_{12}, V^1] = +1V^1$$

- It appears all lowest  $sp(4, \mathbb{R})$  states for 2 particles irreps are of the form

$$T^{2n}|00\rangle|00\rangle \sim V^{2n}|00\rangle|00\rangle \sim V^{2k}T^{2(n-k)}|00\rangle|00\rangle$$

## Two-particle irreps of $sp(4, \mathbb{R})$

- Use  $O(2)$  by constructing the tensors

$$T^1 = \frac{1}{2} \left( a_{11}^\dagger - i a_{21}^\dagger \right), \quad V^1 = \frac{1}{2} \left( a_{12}^\dagger - i a_{22}^\dagger \right).$$

$a_{i\alpha}^\dagger$ : particle number  $i$ , mode number  $\alpha$ .

- Note that

$$[L_{12}, T^1] = +1T^1, \quad [L_{12}, V^1] = +1V^1$$

- It appears all lowest  $sp(4, \mathbb{R})$  states for 2 particles irreps are of the form

$$T^{2n}|00\rangle|00\rangle \sim V^{2n}|00\rangle|00\rangle \sim V^{2k}T^{2(n-k)}|00\rangle|00\rangle$$

- Example:  $|m| = 4$  irrep has lowest state

$$x_{11}^4 - 6x_{11}^2x_{21}^2 + x_{21}^4 \sim |40\rangle|00\rangle - \sqrt{6}|20\rangle|20\rangle + |00\rangle|40\rangle$$

## Two-particle irreps of $sp(4, \mathbb{R})$

- Use  $O(2)$  by constructing the tensors

$$T^1 = \frac{1}{2} \left( a_{11}^\dagger - i a_{21}^\dagger \right), \quad V^1 = \frac{1}{2} \left( a_{12}^\dagger - i a_{22}^\dagger \right).$$

$a_{i\alpha}^\dagger$ : particle number  $i$ , mode number  $\alpha$ .

- Note that

$$[L_{12}, T^1] = +1T^1, \quad [L_{12}, V^1] = +1V^1$$

- It appears all lowest  $sp(4, \mathbb{R})$  states for 2 particles irreps are of the form

$$T^{2n}|00\rangle|00\rangle \sim V^{2n}|00\rangle|00\rangle \sim V^{2k}T^{2(n-k)}|00\rangle|00\rangle$$

- Example:  $|m| = 4$  irrep has lowest state

$$x_{11}^4 - 6x_{11}^2x_{21}^2 + x_{21}^4 \sim |40\rangle|00\rangle - \sqrt{6}|20\rangle|20\rangle + |00\rangle|40\rangle$$

- Compare the  $su(1, 1)$  states at the bottom of  $|m| = 4$ :

$$x_1^4 - 6x_1^2x_2^2 + x_2^4 \sim |4\rangle|0\rangle - \sqrt{6}|2\rangle|2\rangle + |0\rangle|4\rangle$$

# Three-particle irreps of $sp(4, \mathbb{R})$

- We now have

$$A_{ij} = \sum_{k=1}^3 a_{ki}^{\dagger} a_{kj}^{\dagger},$$

$$C_{ij} = \sum_{k=1}^3 \frac{1}{2} \left( a_{ki}^{\dagger} a_{kj} + a_{kj} a_{ki}^{\dagger} \right),$$

$$B_{ij} = A_{ij}^{\dagger}.$$

# Three-particle irreps of $sp(4, \mathbb{R})$

- We now have

$$A_{ij} = \sum_{k=1}^3 a_{ki}^\dagger a_{kj}^\dagger,$$

$$C_{ij} = \sum_{k=1}^3 \frac{1}{2} \left( a_{ki}^\dagger a_{kj} + a_{kj} a_{ki}^\dagger \right),$$

$$B_{ij} = A_{ij}^\dagger.$$

- Use  $O(3)$  by constructing the tensors  $T_M^L$  and  $V_M^L$ :

$$T_{-1}^1 = -\frac{1}{2} \left( a_{11}^\dagger + i a_{21}^\dagger \right), \quad T_0^1 = -\frac{1}{\sqrt{2}} a_{31}^\dagger, \quad T_1^1 = \frac{1}{2} \left( a_{11}^\dagger - i a_{21}^\dagger \right),$$

$$V_{-1}^1 = -\frac{1}{2} \left( a_{12}^\dagger + i a_{22}^\dagger \right), \quad V_0^1 = -\frac{1}{\sqrt{2}} a_{32}^\dagger, \quad V_1^1 = \frac{1}{2} \left( a_{12}^\dagger - i a_{22}^\dagger \right).$$

## Three-particle irreps of $sp(4, \mathbb{R})$

- **Not all** lowest  $sp(4, \mathbb{R})$  states for 3 particles irreps are of the form

$$T^{2n}|00\rangle|00\rangle$$

# Three-particle irreps of $sp(4, \mathbb{R})$

- **Not all** lowest  $sp(4, \mathbb{R})$  states for 3 particles irreps are of the form

$$T^{2n}|00\rangle|00\rangle$$

- Some are:  $L = 2, |m| = 0$  irrep has lowest state

$$x_{11}^2 + x_{21}^2 - 2x_{31}^2 \sim |20\rangle|00\rangle|00\rangle + |00\rangle|20\rangle|00\rangle - 2|00\rangle|00\rangle|20\rangle$$

- Compare the  $su(1, 1)$  states at the bottom of  $L = 2, |m| = 0$ :

$$x_1^2 + x_2^2 - 2x_3^2 \sim |2\rangle|0\rangle|0\rangle + |0\rangle|2\rangle|0\rangle - 2|0\rangle|0\rangle|2\rangle$$



# Three-particle irreps of $sp(4, \mathbb{R})$

- **Not all** lowest  $sp(4, \mathbb{R})$  states for 3 particles irreps are of the form

$$T^{2n}|00\rangle|00\rangle$$

- Some are:  $L = 2, |m| = 0$  irrep has lowest state

$$x_{11}^2 + x_{21}^2 - 2x_{31}^2 \sim |20\rangle|00\rangle|00\rangle + |00\rangle|20\rangle|00\rangle - 2|00\rangle|00\rangle|20\rangle$$

- Compare the  $su(1, 1)$  states at the bottom of  $L = 2, |m| = 0$ :

$$x_1^2 + x_2^2 - 2x_3^2 \sim |2\rangle|0\rangle|0\rangle + |0\rangle|2\rangle|0\rangle - 2|0\rangle|0\rangle|2\rangle$$

- Some aren't:

$$\sum_{mm'} \left\langle \begin{matrix} 2 \\ m \end{matrix}; \begin{matrix} 2 \\ m' \end{matrix} \middle| \begin{matrix} 3 \\ 2 \end{matrix} \right\rangle T_m^2 V_{m'}^2 |00\rangle|00\rangle|00\rangle$$

# Three-particle irreps of $sp(4, \mathbb{R})$

- **Not all** lowest  $sp(4, \mathbb{R})$  states for 3 particles irreps are of the form

$$T^{2n}|00\rangle|00\rangle$$

- Some are:  $L = 2, |m| = 0$  irrep has lowest state

$$x_{11}^2 + x_{21}^2 - 2x_{31}^2 \sim |20\rangle|00\rangle|00\rangle + |00\rangle|20\rangle|00\rangle - 2|00\rangle|00\rangle|20\rangle$$

- Compare the  $su(1, 1)$  states at the bottom of  $L = 2, |m| = 0$ :

$$x_1^2 + x_2^2 - 2x_3^2 \sim |2\rangle|0\rangle|0\rangle + |0\rangle|2\rangle|0\rangle - 2|0\rangle|0\rangle|2\rangle$$

- Some aren't:

$$\sum_{mm'} \left\langle \begin{matrix} 2 \\ m \end{matrix}; \begin{matrix} 2 \\ m' \end{matrix} \middle| \begin{matrix} 3 \\ 2 \end{matrix} \right\rangle T_m^2 V_{m'}^2 |00\rangle|00\rangle|00\rangle$$

- $T^2$  and  $V^2$  are  $L = 2$  tensors under  $O(3)$ , and the combo

$\sum_{mm'} \left\langle \begin{matrix} 2 \\ m \end{matrix}; \begin{matrix} 2 \\ m' \end{matrix} \middle| \begin{matrix} 3 \\ 2 \end{matrix} \right\rangle T_m^2 V_{m'}^2$ , is an “axial tensor” in the sense that it lives in the antisymmetric space of the decomposition  $(L = 2) \otimes (L = 2)$ .

# Three-particle irreps of $sp(4, \mathbb{R})$

- Some don't exist:

$$\sum_{m m'} \left\langle \begin{matrix} 2 \\ m \end{matrix}; \begin{matrix} 2 \\ m' \end{matrix} \mid \begin{matrix} 1 \\ 0 \end{matrix} \right\rangle T_m^2 V_{m'}^2 |00\rangle |00\rangle |00\rangle$$

does not yield product states with only even number of excitations for each particles.

- This is because  $m = 0$  irreps of  $O(2)$  (as a subgroup of  $O(3)$ ) must have even parity but the coupling is antisymmetric in  $T$  and  $V$ .

# Conclusion

- Angular momentum helps in constructing multiple states when combining degenerate representations of  $\mathfrak{su}(1, 1)$ .

# Conclusion

- Angular momentum helps in constructing multiple states when combining degenerate representations of  $\mathfrak{su}(1, 1)$ .
- It is also helpful for 2-particle irreps of  $\mathfrak{sp}(2k, \mathbb{R})$ .

# Conclusion

- Angular momentum helps in constructing multiple states when combining degenerate representations of  $\mathfrak{su}(1, 1)$ .
- It is also helpful for 2-particle irreps of  $\mathfrak{sp}(2k, \mathbb{R})$ .
- Some extra care required for  $n \geq 3$  particles

